

# Sagepath Labs Pvt. Ltd.

Lab Address:- # Plot No. 564 , 1st floor , Buddhanagar , Near Sai Baba Temple Peerzadiguda Boduppal Hyderabad, Telangana. ICMR Reg .No. SAPALAPVLHT (Covid -19)

## LABORATORY TEST REPORT

Name : Mrs. G MADHURI

Sample ID : A0787996

Age/Gender : 50 Years/Female Reg. No : 0312410310009

Referred by : Dr. P S DATATREYA SPP Code : SPL-CV-172

Referring Customer : V CARE MEDICAL DIAGNOSTICS Collected On : 31-Oct-2024 11:18 AM
Primary Sample : Whole Blood Received On : 31-Oct-2024 01:24 PM
Sample Tested In : Whole Blood EDTA Reported On : 31-Oct-2024 03:49 PM

Client Address : Kimtee colony , Gokul Nagar, Tarnaka Report Status : Final Report

HAEMATOLOGY				
Test Name	Results	Units	Biological Reference Interval	
Complete Blood Picture(CBP)				
Maemoglobin (Hb)	<u>11.5</u>	g/dL	12-15	
Haematocrit (HCT)	<u>34.6</u>	%	40-50	
® RBC Count	4.41	10^12/L	3.8-4.8	
(Method: Cell Impedence)  MCV (Method: Calculated)	<u>78</u>	fl	81-101	
⊚ MCH	<u>26.0</u>	pg	27-32	
(Method: Calculated)  MCHC	33.1	g/dL	32.5-34.5	
(Method: Calculated)  RDW-CV (Method: Calculated)	<u>17.5</u>	%	11.6-14.0	
Platelet Count (PLT) (Method: Cell Impedance )	214	10^9/L	150-410	
Total WBC Count (Method: Impedance)	<u>13.4</u>	10^9/L	4.0-10.0	
Differential Leucocyte Count (DC)				
Neutrophils (Method: Cell Impedence)	<u>75</u>	%	40-70 alth Care	
Lymphocytes (Method: Cell Impedence)	20	%	20-40	
Monocytes (Method: Microscopy)	03	%	2-10	
(Method: Microscopy)	02	%	1-6	
Basophils (Method: Microscopy)	00	%	1-2	
(Method: Impedence)	<u>10.05</u>	10^9/L	2.0-7.0	
Mbsolute Lymphocyte Count (Method: Impedence)	2.68	10^9/L	1.0-3.0	
Absolute Monocyte Count     (Method: Calculated)	0.4	10^9/L	0.2-1.0	
Absolute Eosinophils Count     (Method: Calculated)	0.27	10^9/L	0.02-0.5	
Absolute Basophil ICount (Method: Calculated)	0.00	10^9/L	0.0-0.3	
Morphology (Method: PAPs Staining )	Anisocytosis	with Normocy	tic normochromic with Neutrophilic Leucocytosis	









: Serum, Plasma-NaF(R)

# Sagepath Labs Pvt. Ltd.

Lab Address:- # Plot No. 564, 1st floor, Buddhanagar, Near Sai Baba Temple Peerzadiguda Boduppal Hyderabad, Telangana. ICMR Reg. No. SAPALAPVLHT (Covid -19)

: 31-Oct-2024 01:55 PM

## LABORATORY TEST REPORT

Reported On

Name : Mrs. G MADHURI

Sample ID : A0787994, A0787997

Age/Gender : 50 Years/Female Reg. No : 0312410310009

Referred by : Dr. P S DATATREYA SPP Code : SPL-CV-172

Referring Customer : V CARE MEDICAL DIAGNOSTICS Collected On : 31-Oct-2024 11:18 AM Primary Sample : Whole Blood Received On : 31-Oct-2024 01:24 PM

Client Address : Kimtee colony ,Gokul Nagar,Tarnaka Report Status : Final Report

CLINICAL BIOCHEMISTRY				
Test Name	Results	Units	Biological Reference Interval	
Bilirubin(Total) (Method: Diazo)	0.3	mg/dL	0.3-1.2	
Bilirubin (Direct)	0.1	mg/dL	0.0 - 0.3	
Bilirubin (Indirect) (Method: Calculated)	0.2	mg/dL	0.2-1.0	

### **Interpretation:**

Sample Tested In

Bilirubin is a yellowish pigment found in bile, a fluid made by the liver.

Bilirubin is left after these older blood cells are removed. The liver helps break down bilirubin so that it can be removed from the body in the stool. A level of bilirubin in the blood of 2.0 mg/dL can lead to jaundice. Jaundice is a yellow color in the skin, mucus membranes, or eyes.

In newborns, bilirubin level is higher for the first few days of life. Your child's provider must consider the following when deciding whether your baby's bilirubin level is too high:

- How fast the level has been rising
- Whether the baby was born early
- The baby's age

Jaundice can also occur when more red blood cells than normal are broken down. This can be caused by:

- A blood disorder called erythroblastosis fetalis
- A red blood cell disorder called hemolytic anemia
- Transfusion reaction in which red blood cells that were given in a transfusion are destroyed by the person's immune system

Note: DPD(3,5-dichlorophenyldiazonium tetrafluoroborate)

Glucose Random (RBS)	<u>142</u>	mg/dL	70-140
		•	

Interpretation of Plasma Glucose based on ADA guidelines 2018

interpretation of Flashia Glacose based on ADA guidennes 2016				
		2hrsPlasma Glucose(mg/dL)	HbA1c(%)	RBS(mg/dL)
Prediabetes	100-125	140-199	5.7-6.4	NA
Diabetes	> = 126	>= 200	I	>=200(with symptoms)

Reference: Diabetes care 2018:41(suppl.1):S13-S27

- The random blood glucose if it is above 200 mg/dL and the patient has increased thirst, polyuria, and polyphagia, suggests diabetes mellitus.
- As a rule, two-hour glucose samples will reach the fasting level or it will be in the normal range.







Page 2 of 3

DR.VAISHNAVI
MD BIOCHEMISTRY



# Sagepath Labs Pvt. Ltd.

Lab Address:- # Plot No. 564, 1st floor, Buddhanagar, Near Sai Baba Temple Peerzadiguda Boduppal Hyderabad, Telangana. ICMR Reg. No. SAPALAPVLHT (Covid -19)

## LABORATORY TEST REPORT

Name : Mrs. G MADHURI

Sample ID : A0787994, A0787997

Age/Gender : 50 Years/Female Reg. No : 0312410310009

Referred by : Dr. P S DATATREYA SPP Code : SPL-CV-172

Referring Customer : V CARE MEDICAL DIAGNOSTICS Collected On : 31-Oct-2024 11:18 AM Primary Sample : Whole Blood Received On : 31-Oct-2024 01:24 PM

Sample Tested In : Serum, Plasma-NaF(R) Reported On : 31-Oct-2024 01:55 PM

Client Address : Kimtee colony ,Gokul Nagar,Tarnaka Report Status : Final Report

CLINICAL BIOCHEMISTRY			
Test Name	Results	Units	Biological Reference Interval
Creatinine (Method: Jaffes Kinetic)	0.62	mg/dL	0.60-1.10

### **Interpretation:**

- This test is done to see how well your kidneys are working. Creatinine is a chemical waste product of creatine. Creatine is a chemical made by the body and is used to supply energy mainly to muscles.
- A higher than normal level may be due to:
- Renal diseases and insufficiency with decreased glomerular filtration, urinary tract obstruction, reduced renal blood flow including congestive heart failure, shock, and dehydration; rhabdomyolysis can cause elevated serum creatinine.
- A lower than normal level may be due to:
- Small stature, debilitation, decreased muscle mass; some complex cases of severe hepatic disease can cause low serum creatinine levels. In advanced liver disease, low creatinine may result from decreased hepatic production of creatinine and inadequate dietary protein as well as reduced musle mass.

\*\*\* End Of Report \*\*\*

Excellence In Health Care







Page 3 of 3

DR. VAISHNAVI
MD BIOCHEMISTRY