










LABORATORY TEST REPORT

Name	: Mrs. RENUKA		
Sample ID	: A0788101		
Age/Gender	: 34 Years/Female	Reg. No	: 0312411040013
Referred by	: Dr. KRISHNA YAJI	SPP Code	: SPL-CV-172
Referring Customer	: V CARE MEDICAL DIAGNOSTICS	Collected On	: 04-Nov-2024 10:59 AM
Primary Sample	: Whole Blood	Received On	: 04-Nov-2024 12:50 PM
Sample Tested In	: Whole Blood EDTA	Reported On	: 04-Nov-2024 01:10 PM
Client Address	: Kimtee colony ,Gokul Nagar,Tarnaka	Report Status	: Final Report












HAEMATOLOGY

Test Name	Results	Units	Biological Reference Interval
-----------	---------	-------	-------------------------------

Complete Blood Picture(CBP)

 Haemoglobin (Hb) (Method: Cymeth Method)	6.6	g/dL	12-15
 Haematocrit (HCT) (Method: Calculated)	23.9	%	40-50
 RBC Count (Method: Cell Impedance)	3.91	10 ¹² /L	3.8-4.8
 MCV (Method: Calculated)	61	fl	81-101
 MCH (Method: Calculated)	16.9	pg	27-32
 MCHC (Method: Calculated)	27.7	g/dL	32.5-34.5
 RDW-CV (Method: Calculated)	18.9	%	11.6-14.0
 Platelet Count (PLT) (Method: Cell Impedance)	272	10 ⁹ /L	150-410
 Total WBC Count (Method: Impedance)	8.7	10 ⁹ /L	4.0-10.0

Differential Leucocyte Count (DC)

 Neutrophils (Method: Cell Impedance)	67	%	40-70
 Lymphocytes (Method: Cell Impedance)	27	%	20-40
 Monocytes (Method: Microscopy)	04	%	2-10
 Eosinophils (Method: Microscopy)	02	%	1-6
 Basophils (Method: Microscopy)	00	%	1-2
 Absolute Neutrophils Count (Method: Impedance)	5.83	10 ⁹ /L	2.0-7.0
 Absolute Lymphocyte Count (Method: Impedance)	2.35	10 ⁹ /L	1.0-3.0
 Absolute Monocyte Count (Method: Calculated)	0.35	10 ⁹ /L	0.2-1.0
 Absolute Eosinophils Count (Method: Calculated)	0.17	10 ⁹ /L	0.02-0.5
 Absolute Basophil ICount (Method: Calculated)	0.00	10 ⁹ /L	0.0-0.3

Morphology

(Method: PAPs Staining)

Anisocytosis with Severe Microcytic hypochromic anemia



LABORATORY TEST REPORT

Name	: Mrs. RENUKA		
Sample ID	: A0788102		
Age/Gender	: 34 Years/Female	Reg. No	: 0312411040013
Referred by	: Dr. KRISHNA YAJI	SPP Code	: SPL-CV-172
Referring Customer	: V CARE MEDICAL DIAGNOSTICS	Collected On	: 04-Nov-2024 10:59 AM
Primary Sample	:	Received On	: 04-Nov-2024 12:50 PM
Sample Tested In	: Urine	Reported On	: 04-Nov-2024 02:28 PM
Client Address	: Kimtee colony ,Gokul Nagar,Tarnaka	Report Status	: Final Report


CLINICAL PATHOLOGY

Test Name	Results	Units	Biological Reference Interval
-----------	---------	-------	-------------------------------

Complete Urine Analysis (CUE)
Physical Examination

Colour	Pale Yellow	Straw to light amber
Appearance	HAZY	Clear

Chemical Examination

Glucose <small>(Method: Strip Reflectance)</small>	Negative	Negative
Protein <small>(Method: Strip Reflectance)</small>	(+)	Negative
Bilirubin (Bile) <small>(Method: Strip Reflectance)</small>	Negative	Negative
Urobilinogen <small>(Method: Ehrlichs reagent)</small>	Negative	Negative
Ketone Bodies <small>(Method: Strip Reflectance)</small>	Negative	Negative
Specific Gravity <small>(Method: Strip Reflectance)</small>	1.025	1.000 - 1.030
Blood <small>(Method: Strip Reflectance)</small>	Negative	Negative
Reaction (pH) <small>(Method: Reagent Strip Reflectance)</small>	6.0	5.0 - 8.5
Nitrites <small>(Method: Strip Reflectance)</small>	Negative	Negative
Leukocyte esterase <small>(Method: Reagent Strip Reflectance)</small>	Negative	Negative

Microscopic Examination (Microscopy)

PUS(WBC) Cells <small>(Method: Microscopy)</small>	03-04	/hpf	00-05
R.B.C. <small>(Method: Microscopic)</small>	Nil	/hpf	Nil
Epithelial Cells <small>(Method: Microscopic)</small>	02-03	/hpf	00-05
Casts <small>(Method: Microscopic)</small>	Absent		Absent
Crystals <small>(Method: Microscopic)</small>	Absent		Absent
Bacteria	Nil		Nil
Budding Yeast Cells <small>(Method: Microscopy)</small>	Nil		Absent

Comments :Urine analysis is one of the most useful laboratory tests as it identifies a wide range of medical conditions including renal damage, urinary tract infections,diabetes, hypertension and drug toxicity.


 Page 2 of 3
 Swarnabala - M
 DR.SWARNA BALA
 MD PATHOLOGY

LABORATORY TEST REPORT

Name	: Mrs. RENUKA		
Sample ID	: A0788054		
Age/Gender	: 34 Years/Female	Reg. No	: 0312411040013
Referred by	: Dr. KRISHNA YAJI	SPP Code	: SPL-CV-172
Referring Customer	: V CARE MEDICAL DIAGNOSTICS	Collected On	: 04-Nov-2024 10:59 AM
Primary Sample	: Whole Blood	Received On	: 04-Nov-2024 12:50 PM
Sample Tested In	: Serum	Reported On	: 04-Nov-2024 02:09 PM
Client Address	: Kimtee colony ,Gokul Nagar,Tarnaka	Report Status	: Final Report


CLINICAL BIOCHEMISTRY

Test Name	Results	Units	Biological Reference Interval
-----------	---------	-------	-------------------------------


TSH -Thyroid Stimulating Hormone
1.34
μIU/mL
0.35-5.5

(Method: CLIA)

Pregnancy & Cord Blood

TSH (Thyroid Stimulating Hormone (μIU/mL))	
First Trimester	: 0.24-2.99
Second Trimester	: 0.46-2.95
Third Trimester	: 0.43-2.78
Cord Blood	: 2.3-13.2

- TSH is synthesized and secreted by the anterior pituitary in response to a negative feedback mechanism involving concentrations of FT3 (free T3) and FT4 (free T4). Additionally, the hypothalamic tripeptide, thyrotropin-releasing hormone (TRH), directly stimulates TSH production.
- TSH interacts with specific cell receptors on the thyroid cell surface and exerts two main actions. The first action is to stimulate cell reproduction and hypertrophy. Secondly, TSH stimulates the thyroid gland to synthesize and secrete T3 and T4
- The ability to quantitate circulating levels of TSH is important in evaluating thyroid function. It is especially useful in the differential diagnosis of primary (thyroid) from secondary (pituitary) and tertiary (hypothalamus) hypothyroidism. In primary hypothyroidism, TSH levels are significantly elevated, while in secondary and tertiary hypothyroidism, TSH levels are low
- TRH stimulation differentiates secondary and tertiary hypothyroidism by observing the change in patient TSH levels. Typically, the TSH response to TRH stimulation is absent in cases of secondary hypothyroidism, and normal to exaggerated in tertiary hypothyroidism
- Historically, TRH stimulation has been used to confirm primary hyperthyroidism, indicated by elevated T3 and T4 levels and low or undetectable TSH levels. TSH assays with increased sensitivity and specificity provide a primary diagnostic tool to differentiate hyperthyroid from euthyroid patients.

*** End Of Report ***




DR. VAISHNAVI
MD BIOCHEMISTRY

Page 3 of 3