



Sagepath Labs Pvt. Ltd.

Lab Address:- # Plot No. 564, 1st floor, Buddhanagar, Near Sai Baba Temple Peerzadiguda Boduppal Hyderabad, Telangana. ICMR Reg. No. SAPALAPVLHT (Covid -19)

LABORATORY TEST REPORT

Name : Mrs. CHINNAPPA MERIBAH PAUL Sample ID : 24202322, 24202323, 24202321

Age/Gender : 54 Years/Female Reg. No : 0312411090004

Referred by : Dr. SELF SPP Code : SPL-CV-172
Referring Customer : V CARE MEDICAL DIAGNOSTICS Collected On : 09-Nov-2024 09:00 AM

Primary Sample : Whole Blood : 09-Nov-2024 01:03 PM Sample Tested In : Plasma-NaF(F), Plasma-NaF(PP), Reported On : 09-Nov-2024 02:29 PM

Client Address : Kimtee colony ,Gokul Nagar,Tarnaka Report Status : Final Report

CLINICAL BIOCHEMISTRY

Test Name Results Units Biological Reference Interval

Glucose Fasting (F) <u>108</u> mg/dL 70-100

Interpretation of Plasma Glucose based on ADA guidelines 2018

Diagnosis	FastingPlasma Glucose(mg/dL)	2hrsPlasma Glucose(mg/dL)	HbA1c(%)	RBS(mg/dL)	
Prediabetes	100-125	140-199	5.7-6.4	NA	
Diabetes	>= 126	>= 200	> = 6.5	>=200(with symptoms)	

Reference: Diabetes care 2018:41(suppl.1):S13-S27

Glucose Post Prandial (PP) 139 mg/dL 70-140

Interpretation of Plasma Glucose based on ADA guidelines 2018

Diagnosis	FastingPlasma Glucose(mg/dL)	2hrsPlasma Glucose(mg/dL)	HbA1c(%)	RBS(mg/dL)
Prediabetes	100-125	140-199	5.7-6.4	NA
Diabetes	> = 126	>= 200	>= 6.5	>=200(with symptoms)


Reference: Diabetes care 2018:41(suppl.1):S13-S27

- Postprandial glucose level is a screening test for Diabetes Mellitus
- $\bullet \quad \text{If glucose level is} > 140 \text{ mg/dL and} < 200 \text{ mg/dL}, \text{ then GTT (glucose tolerance test) is advised.}$
- If level after 2 hours = >200 mg/dL diabetes mellitus is confirmed.
- Advise HbA1c for further evaluation.











Sagepath Labs Pvt. Ltd.

Lab Address:- # Plot No. 564, 1st floor, Buddhanagar, Near Sai Baba Temple Peerzadiguda Boduppal Hyderabad, Telangana. ICMR Reg. No. SAPALAPVLHT (Covid -19)

LABORATORY TEST REPORT

Name : Mrs. CHINNAPPA MERIBAH PAUL Sample ID : 24202322, 24202323, 24202321

Sample 1D : 24202322, 24202323, 2420232 Age/Gender : 54 Years/Female

Referred by : Dr. SELF

Referring Customer : V CARE MEDICAL DIAGNOSTICS
Primary Sample : Whole Blood

Sample Tested In : Plasma-NaF(F), Plasma-NaF(PP),

Client Address : Kimtee colony ,Gokul Nagar,Tarnaka

Reg. No : 0312411090004

SPP Code : SPL-CV-172

Collected On : 09-Nov-2024 09:00 AM Received On : 09-Nov-2024 01:03 PM

Reported On : 09-Nov-2024 02:29 PM

Report Status : Final Report

CLINICAL BIOCHEMISTRY				
Test Name	Results	Units	Biological Reference Interval	
TSH -Thyroid Stimulating Hormone	<u>33.36</u>	μIU/mL	0.35-5.5	

Pregnancy & Cord Blood				
		TSH (Thyroid Stimulating Hormone (μIU/mL)		
First Trimester	: 0.24-2.99			
Second Trimester	Second Trimester: 0.46-2.95			
Third Trimester	: 0.43-2.78			
Cord Blood	: 2.3-13.2			

- TSH is synthesized and secreted by the anterior pituitary in response to a negative feedback mechanism involving concentrations of FT3 (free T3) and FT4 (free T4). Additionally, the hypothalamic tripeptide, thyrotropin-releasing hormone (TRH), directly stimulates TSH production.
- TSH interacts with specific cell receptors on the thyroid cell surface and exerts two main actions. The first action is to stimulate cell reproduction and hypertrophy. Secondly, TSH stimulates the thyroid gland to synthesize and secrete T3 and T4
- The ability to quantitate circulating levels of TSH is important in evaluating thyroid function. It is especially useful in the differential diagnosis of primary (thyroid) from secondary (pituitary) and tertiary (hypothalamus) hypothyroidism. In primary hypothyroidism, TSH levels are significantly elevated, while in secondary and tertiary hypothyroidism, TSH levels are low
- TRH stimulation differentiates secondary and tertiary hypothyroidism by observing the change in patient TSH levels. Typically, the TSH response to TRH stimulation is absent in cases of secondary hypothyroidism, and normal to exaggerated in tertiary hypothyroidism
- Historically, TRH stimulation has been used to confirm primary hyperthyroidism, indicated by elevated T3 and T4 levels and low or undetectable TSH levels. TSH assays with increased sensitivity and specificity provide a primary diagnostic tool to differentiate hyperthyroid from euthyroid patients.

*** End Of Report ***







Page 2 of 2

DR. VAISHNAVI
MD BIOCHEMISTRY