

ITDOSE INFOSYSTEMS PVT. LTD.

Sagepath Labs Pvt. Ltd.

Lab Address:- # Plot No. 564 , 1st floor , Buddhanagar , Near Sai Baba Temple Peerzadiguda Boduppal Hyderabad, Telangana. ICMR Reg .No. SAPALAPVLHT (Covid -19)

LABORATORY TEST REPORT

Name	: Mrs. G MADHURI		
Sample ID	: 24202502		
Age/Gender	: 49 Years/Female	Reg. No	: 0312411130012
Referred by	: Dr. P SATYA DATTATREYA	SPP Code	: SPL-CV-172
Referring Customer	: V CARE MEDICAL DIAGNOSTICS	Collected On	: 13-Nov-2024 11:22 AM
Primary Sample	: Whole Blood	Received On	: 13-Nov-2024 12:54 PM
Sample Tested In	: Whole Blood EDTA	Reported On	: 13-Nov-2024 02:02 PM
Client Address	: Kimtee colony ,Gokul Nagar,Tarnaka	Report Status	: Final Report

HAEMATOLOGY						
Test Name	Results	Units	Biological Reference Interval			
Complete Blood Picture(CBP)						
Haemoglobin (Hb) (Methad: Cynmeth Methad)	<u>10.0</u>	g/dL	12-15			
Haematocrit (HCT)	<u>30.4</u>	%	40-50			
RBC Count Method: Cell Impedence)	<u>3.59</u>	10^12/L	3.8-4.8			
(Method: Calculated)	84	fl	81-101			
(Method: Calculated)	27.9	pg	27-32			
(Method: Calculated)	33.0	g/dL	32.5-34.5			
RDW-CV (Nethod: Calculated)	<u>18.1</u>	%	11.6-14.0			
Platelet Count (PLT) Count (PLT) Plated: Cell Impedance)	196	10^9/L	150-410			
Total WBC Count Method: Impedance)	9.9	10^9/L	4.0-10.0			
Differential Leucocyte Count (DC)						
Neutrophils (Method: Cell Impedence)	70 Ce	%	40-70 alth Care			
(Method: Cell Impedence)	20	%	20-40			
Monocytes	06	%	2-10			
Eosinophils (Method: Microscopy)	04	%	1-6			
Basophils (Method: Microscopy)	00	%	1-2			
	6.93	10^9/L	2.0-7.0			
Absolute Lymphocyte Count Method: Impedence)	1.98	10^9/L	1.0-3.0			
Absolute Monocyte Count Monocyte Count	0.59	10^9/L	0.2-1.0			
	0.4	10^9/L	0.02-0.5			
Absolute Basophil ICount Method: Calculated)	0.00	10^9/L	0.0-0.3			
Morphology (Method: PAPS Staining)	Anisocytosis	with Normocyt	tic normochromic anemia			





Page 1 of 3 Swarnabala.M DR.SWARNA BALA MD PATHOLOGY

Note : This report is subject to the terms and conditions overleaf. Partial Reproduction of this report is not Permitted



Sagepath Labs Pvt. Ltd.

Lab Address:- # Plot No. 564 , 1st floor , Buddhanagar , Near Sai Baba Temple Peerzadiguda Boduppal Hyderabad, Telangana. ICMR Reg .No. SAPALAPVLHT (Covid -19)

LABORATORY TEST REPORT

I	Name	: Mrs. G MADHURI		
I	Sample ID	: 24202501, 24202483		
I	Age/Gender	: 49 Years/Female	Reg. No	: 0312411130012
I	Referred by	: Dr. P SATYA DATTATREYA	SPP Code	: SPL-CV-172
I	Referring Customer	: V CARE MEDICAL DIAGNOSTICS	Collected On	: 13-Nov-2024 11:22 AM
I	Primary Sample	: Whole Blood	Received On	: 13-Nov-2024 12:48 PM
I	Sample Tested In	: Serum, Plasma-NaF(R)	Reported On	: 13-Nov-2024 02:39 PM
I	Client Address	: Kimtee colony ,Gokul Nagar,Tarnaka	Report Status	: Final Report

CLINICAL BIOCHEMISTRY				
Test Name	Results	Units	Biological Reference Interval	
Bilirubin(Total)	0.5	mg/dL	0.3-1.2	
Bilirubin (Direct)	0.1	mg/dL	0.0 - 0.3	
	0.4	mg/dL	0.2-1.0	

Interpretation:

Bilirubin is a yellowish pigment found in bile, a fluid made by the liver.

Bilirubin is left after these older blood cells are removed. The liver helps break down bilirubin so that it can be removed from the body in the stool. A level of bilirubin in the blood of 2.0 mg/dL can lead to jaundice. Jaundice is a yellow color in the skin, mucus membranes, or eyes.

In newborns, bilirubin level is higher for the first few days of life. Your child's provider must consider the following when deciding whether your baby's bilirubin level is too high:

- How fast the level has been rising
- Whether the baby was born early
- The baby's age

Jaundice can also occur when more red blood cells than normal are broken down. This can be caused by:

- A blood disorder called erythroblastosis fetalis
- A red blood cell disorder called hemolytic anemia
- Transfusion reaction in which red blood cells that were given in a transfusion are destroyed by the person's immune system

Note: DPD(3,5-dichlorophenyldiazonium tetrafluoroborate)

Glucose Random (RBS)	124	mg/dL	70-140	
(Method: Heyekingso (HK))		-		

Interpretation of Plasma Glucose based on ADA guidelines 2018							
Diagnosis	FastingPlasma Glucose(mg/dL)	2hrsPlasma Glucose(mg/dL)	HbA1c(%)	RBS(mg/dL)			
Prediabetes	100-125	140-199	5.7-6.4	NA			
Diabetes	> = 126	> = 200		>=200(with symptoms)			

Reference: Diabetes care 2018:41(suppl.1):S13-S27

• The random blood glucose if it is above 200 mg/dL and the patient has increased thirst, polyuria, and polyphagia, suggests diabetes mellitus.

• As a rule, two-hour glucose samples will reach the fasting level or it will be in the normal range.







Page 2 of 3 BIOCHEMISTRY

Note : This report is subject to the terms and conditions overleaf. Partial Reproduction of this report is not Permitted



Sagepath Labs Pvt. Ltd.

Lab Address:- # Plot No. 564 , 1st floor , Buddhanagar , Near Sai Baba Temple Peerzadiguda Boduppal Hyderabad, Telangana. ICMR Reg .No. SAPALAPVLHT (Covid -19)

LABORATORY TEST REPORT

Name	: Mrs. G MADHURI		
Sample ID	: 24202501, 24202483		
Age/Gender	: 49 Years/Female	Reg. No	: 0312411130012
Referred by	: Dr. P SATYA DATTATREYA	SPP Code	: SPL-CV-172
Referring Customer	: V CARE MEDICAL DIAGNOSTICS	Collected On	: 13-Nov-2024 11:22 AM
Primary Sample	: Whole Blood	Received On	: 13-Nov-2024 12:48 PM
Sample Tested In	: Serum, Plasma-NaF(R)	Reported On	: 13-Nov-2024 02:39 PM
Client Address	: Kimtee colony ,Gokul Nagar,Tarnaka	Report Status	: Final Report

CLINICAL BIOCHEMISTRY				
Test Name	Results	Units	Biological Reference Interval	
	0.63	mg/dL	0.60-1.10	

Interpretation:

- This test is done to see how well your kidneys are working. Creatinine is a chemical waste product of creatine. Creatine is a chemical made by the body and is used to supply energy mainly to muscles.
- A higher than normal level may be due to:
- Renal diseases and insufficiency with decreased glomerular filtration, urinary tract obstruction, reduced renal blood flow including congestive heart failure, shock, and dehydration; rhabdomyolysis can cause elevated serum creatinine.
- A lower than normal level may be due to:
- Small stature, debilitation, decreased muscle mass; some complex cases of severe hepatic disease can cause low serum creatinine levels. In advanced liver disease, low creatinine may result from decreased hepatic production of creatinine and inadequate dietary protein as well as reduced musle mass.

*** End Of Report ***







Page 3 of 3 MD BIOCHEMISTRY