

Sagepath Labs Pvt. Ltd.

Lab Address:- # Plot No. 564 , 1st floor , Buddhanagar , Near Sai Baba Temple Peerzadiguda Boduppal Hyderabad, Telangana. ICMR Reg .No. SAPALAPVLHT (Covid -19)

LABORATORY TEST REPORT

Name	: Mrs. J LAVNYA		
Sample ID	: A1307642, A1307644, A1307641		
Age/Gender	: 38 Years/Female	Reg. No	: 0312411160007
Referred by	: Dr. SELF	SPP Code	: SPL-CV-172
Referring Customer	: V CARE MEDICAL DIAGNOSTICS	Collected On	: 16-Nov-2024 10:34 AM
Primary Sample	: Whole Blood	Received On	: 16-Nov-2024 12:38 PM
Sample Tested In	: Plasma-NaF(F), Plasma-NaF(PP),	Reported On	: 16-Nov-2024 02:29 PM
Client Address	: Kimtee colony ,Gokul Nagar,Tarnaka	Report Status	: Final Report

est Name		Results	Units	Biological Refer	ence Interva
Glucose Fa (Method: Hexokinase)		<u>112</u>	mg/dL	70-100	
nterpretation of I	Plasma Glucose based on ADA guidelines	\$ 2018			
Diagnosis	FastingPlasma Glucose(mg/dL)	2hrsPlasma Glucose(m	g/dL) HbA1c	%) RBS(mg/dL)	
Prediabetes	100-125	140-199	5.7-6.4	NA	
			1	000/	c)
Reference: Dial	> = 126 betes care 2018:41(suppl.1):S13-S2		> = 6.9		5)
Glucose Po (Method: Hexokinase	betes care 2018:41(suppl.1):S13-S2 ost Prandial (PP)	7 120	> = 6.9	>=200(with sympton 70-140	s)
Reference: Dial Glucose Po (Method: Hexokinase Interpretation of	betes care 2018:41(suppl.1):S13-S2 ost Prandial (PP)	7 120	mg/dL	70-140	•)
Reference: Dial Glucose Po (Method: Hexokinase Interpretation of	betes care 2018:41(suppl.1):S13-S2 Ost Prandial (PP) Plasma Glucose based on ADA guideline	7 120 \$ 2018	mg/dL	70-140	s)

• If glucose level is >140 mg/dL and <200 mg/dL, then GTT (glucose tolerance test) is advised.

• If level after 2 hours = >200 mg/dL diabetes mellitus is confirmed.

• Advise HbA1c for further evaluation.

ITDOSE INFOSYSTEMS PVT. LTD.









Sagepath Labs Pvt. Ltd.

Lab Address:- # Plot No. 564 , 1st floor , Buddhanagar , Near Sai Baba Temple Peerzadiguda Boduppal Hyderabad, Telangana. ICMR Reg .No. SAPALAPVLHT (Covid -19)

LABORATORY TEST REPORT

Name	: Mrs. J LAVNYA		
Sample ID	: A1307642, A1307644, A1307641		
Age/Gender	: 38 Years/Female	Reg. No	: 0312411160007
Referred by	: Dr. SELF	SPP Code	: SPL-CV-172
Referring Customer	: V CARE MEDICAL DIAGNOSTICS	Collected On	: 16-Nov-2024 10:34 AM
Primary Sample	: Whole Blood	Received On	: 16-Nov-2024 12:38 PM
Sample Tested In	: Plasma-NaF(F), Plasma-NaF(PP),	Reported On	: 16-Nov-2024 02:29 PM
Client Address	: Kimtee colony ,Gokul Nagar,Tarnaka	Report Status	: Final Report

CLINICAL BIOCHEMISTRY				
Test Name	Results	Units	Biological Reference Interval	
TSH -Thyroid Stimulating Hormone	4.37	µIU/mL	0.35-5.5	

Pregnancy & Con	rd Blood	
		TSH (Thyroid Stimulating Hormone (µIU/mL)
First Trimester	: 0.24-2.99	
Second Trimester	: 0.46-2.95	
Third Trimester	: 0.43-2.78	
Cord Blood	: 2.3-13.2	

• TSH is synthesized and secreted by the anterior pituitary in response to a negative feedback mechanism involving concentrations of FT3 (free T3) and FT4 (free T4). Additionally, the hypothalamic tripeptide, thyrotropin-releasing hormone (TRH), directly stimulates TSH production.

- TSH interacts with specific cell receptors on the thyroid cell surface and exerts two main actions. The first action is to stimulate cell reproduction and hypertrophy. Secondly, TSH stimulates the thyroid gland to synthesize and secrete T3 and T4
- The ability to quantitate circulating levels of TSH is important in evaluating thyroid function. It is especially useful in the differential diagnosis of primary (thyroid) from secondary (pituitary) and tertiary (hypothalamus) hypothyroidism. In primary hypothyroidism, TSH levels are significantly elevated, while in secondary and tertiary hypothyroidism, TSH levels are low
- TRH stimulation differentiates secondary and tertiary hypothyroidism by observing the change in patient TSH levels. Typically, the TSH response to TRH stimulation is absent in cases of secondary hypothyroidism, and normal to exaggerated in tertiary hypothyroidism
- Historically, TRH stimulation has been used to confirm primary hyperthyroidism, indicated by elevated T3 and T4 levels and low or undetectable TSH levels. TSH assays with increased sensitivity and specificity provide a primary diagnostic tool to differentiate hyperthyroid from euthyroid patients.

*** End Of Report ***







