

**LABORATORY TEST REPORT**

Name	: Mr. KONDAM RAJU		
Sample ID	: A1308658, A1308660		
Age/Gender	: 44 Years/Male	Reg. No	: 0312412260005
Referred by	: Dr. SELF	SPP Code	: SPL-CV-172
Referring Customer	: V CARE MEDICAL DIAGNOSTICS	Collected On	: 26-Dec-2024 11:00 AM
Primary Sample	: Whole Blood	Received On	: 26-Dec-2024 03:05 PM
Sample Tested In	: Plasma-NaF(F), Plasma-NaF(PP)	Reported On	: 26-Dec-2024 03:43 PM
Client Address	: Kimtee colony ,Gokul Nagar,Tarnaka	Report Status	: Final Report


**CLINICAL BIOCHEMISTRY**
**GLUCOSE POST PRANDIAL (PP)**

Test Name	Results	Units	Biological Reference Interval
-----------	---------	-------	-------------------------------

Glucose Fasting (F) 99 mg/dL 70-100  
 (Method: Hexokinase)

Interpretation of Plasma Glucose based on ADA guidelines 2018

Diagnosis	Fasting Plasma Glucose(mg/dL)	2hrs Plasma Glucose(mg/dL)	HbA1c(%)	RBS(mg/dL)
Prediabetes	100-125	140-199	5.7-6.4	NA
Diabetes	> = 126	> = 200	> = 6.5	>=200(with symptoms)

Reference: Diabetes care 2018;41(suppl.1):S13-S27

Glucose Post Prandial (PP) 108 mg/dL 70-140  
 (Method: Hexokinase (HK))

Interpretation of Plasma Glucose based on ADA guidelines 2018

Diagnosis	Fasting Plasma Glucose(mg/dL)	2hrs Plasma Glucose(mg/dL)	HbA1c(%)	RBS(mg/dL)
Prediabetes	100-125	140-199	5.7-6.4	NA
Diabetes	> = 126	> = 200	> = 6.5	>=200(with symptoms)

Reference: Diabetes care 2018;41(suppl.1):S13-S27

- Postprandial glucose level is a screening test for Diabetes Mellitus
- If glucose level is >140 mg/dL and <200 mg/dL, then GTT (glucose tolerance test) is advised.
- If level after 2 hours = >200 mg/dL diabetes mellitus is confirmed.
- Advise HbA1c for further evaluation.

\*\*\* End Of Report \*\*\*



*Dr. Vaishnavi*  
**DR. VAISHNAVI**  
**MD BIOCHEMISTRY**

Page 1 of 3

**LABORATORY TEST REPORT**

Name	: Mr. KONDAM RAJU		
Sample ID	: A1308657		
Age/Gender	: 44 Years/Male	Reg. No	: 0312412260005
Referred by	: Dr. SELF	SPP Code	: SPL-CV-172
Referring Customer	: V CARE MEDICAL DIAGNOSTICS	Collected On	: 26-Dec-2024 11:00 AM
Primary Sample	: Whole Blood	Received On	: 26-Dec-2024 12:52 PM
Sample Tested In	: Serum	Reported On	: 26-Dec-2024 04:27 PM
Client Address	: Kimtee colony ,Gokul Nagar,Tarnaka	Report Status	: Final Report


**CLINICAL BIOCHEMISTRY**

Test Name	Results	Units	Biological Reference Interval
-----------	---------	-------	-------------------------------

**Thyroid Profile-I(TFT)**

 <b>T3 (Triiodothyronine)</b> <small>(Method: CLIA)</small>	132.65	ng/dL	70-204
 <b>T4 (Thyroxine)</b> <small>(Method: CLIA)</small>	9.0	µg/dL	3.2-12.6
 <b>TSH -Thyroid Stimulating Hormone</b> <small>(Method: CLIA)</small>	0.65	µIU/mL	0.35-5.5

**Pregnancy & Cord Blood**

<b>T3 (Triiodothyronine):</b>	<b>T4 (Thyroxine)</b>	<b>TSH (Thyroid Stimulating Hormone)</b>
First Trimester : 81-190 ng/dL	15 to 40 weeks:9.1-14.0 µg/dL	First Trimester : 0.24-2.99 µIU/mL
Second&Third Trimester :100-260 ng/dL		Second Trimester: 0.46-2.95 µIU/mL
		Third Trimester : 0.43-2.78 µIU/mL
Cord Blood: 30-70 ng/dL	Cord Blood: 7.4-13.0 µg/dL	Cord Blood: : 2.3-13.2 µIU/mL

**Interpretation:**

- Thyroid gland is a butterfly-shaped endocrine gland that is normally located in the lower front of the neck. The thyroid's job is to make thyroid hormones, which are secreted into the blood and then carried to every tissue in the body. Thyroid hormones help the body use energy, stay warm and keep the brain, heart, muscles, and other organs working as they should.
- Thyroid produces two major hormones: triiodothyronine (T3) and thyroxine (T4). If thyroid gland doesn't produce enough of these hormones, you may experience symptoms such as weight gain, lack of energy, and depression. This condition is called hypothyroidism.
- Thyroid gland produces too many hormones, you may experience weight loss, high levels of anxiety, tremors, and a sense of being on a high. This is called hyperthyroidism.
- TSH interacts with specific cell receptors on the thyroid cell surface and exerts two main actions. The first action is to stimulate cell reproduction and hypertrophy. Secondly, TSH stimulates the thyroid gland to synthesize and secrete T3 and T4.
- The ability to quantitate circulating levels of TSH is important in evaluating thyroid function. It is especially useful in the differential diagnosis of primary (thyroid) from secondary (pituitary) and tertiary (hypothalamus) hypothyroidism. In primary hypothyroidism, TSH levels are significantly elevated, while in secondary and tertiary hypothyroidism, TSH levels are low.

\*\*\* End Of Report \*\*\*



DR. VAISHNAVI  
MD BIOCHEMISTRY

Page 2 of 3

**LABORATORY TEST REPORT**

Name	: Mr. KONDAM RAJU	Reg. No	: 0312412260005
Sample ID	: A1308657	SPP Code	: SPL-CV-172
Age/Gender	: 44 Years/Male	Collected On	: 26-Dec-2024 11:00 AM
Referred by	: Dr. SELF	Received On	: 26-Dec-2024 12:52 PM
Referring Customer	: V CARE MEDICAL DIAGNOSTICS	Reported On	: 26-Dec-2024 07:29 PM
Primary Sample	: Whole Blood	Report Status	: Final Report
Sample Tested In	: Serum		
Client Address	: Kimtee colony ,Gokul Nagar,Tarnaka		




**IMMUNOLOGY & SEROLOGY**

Test Name	Results	Units	Biological Reference Interval
HIV (1 & 2) Antibody (Method: ELISA)	0.33	S/Co	< 1.00 : Negative > 1.00 : Positive

**Interpretation**

- Non Reactive result implies that antibodies to HIV 1 / 2 have not been detected in the sample. This means the patient has either not been exposed to HIV 1 / 2 infection or the sample has been tested during the "window phase" i.e. before the development of detectable levels of antibodies. Hence a Non Reactive result does not exclude the possibility of exposure or infection with HIV 1 / 2.
- Pre and Post test counseling to be done by the concerned referring doctor. The sensitivity and specificity of this test has been determined by National HIV Reference Centers of Govt. of India and WHO collaborating Centers, using various other test panels. "
- Reactive samples by ELISA Method are confirmed by 2 other supplemental tests for confirm of HIV infection as per NACO guidelines.
- All patients' reports indeterminate should be repeated with a second sample taken 14-28 days. In case the serological results continue to be indeterminate the sample should be subject to western blot for confirmation.



\*\*\* End Of Report \*\*\*



Page 3 of 3

*[Signature]*

**DR. RUTURAJ MANIKLAL KOLHAPURE**  
MD, MICROBIOLOGIST