










LABORATORY TEST REPORT

Name	: Miss. POOJA		
Sample ID	: A1309185		
Age/Gender	: 17 Years/Female	Reg. No	: 0312501160078
Referred by	: Dr. Nivedita Ashrit MD (Obs/Gyn)	SPP Code	: SPL-CV-172
Referring Customer	: V CARE MEDICAL DIAGNOSTICS	Collected On	: 16-Jan-2025 08:08 PM
Primary Sample	: Whole Blood	Received On	: 16-Jan-2025 10:48 PM
Sample Tested In	: Whole Blood EDTA	Reported On	: 16-Jan-2025 10:59 PM
Client Address	: Kimtee colony ,Gokul Nagar,Tarnaka	Report Status	: Final Report












HAEMATOLOGY

Test Name	Results	Units	Biological Reference Interval
-----------	---------	-------	-------------------------------

Complete Blood Picture(CBP)

 Haemoglobin (Hb) <small>(Method: Cymeth Method)</small>	11.9	g/dL	12-15
 Haematocrit (HCT) <small>(Method: Calculated)</small>	41.5	%	40-50
 RBC Count <small>(Method: Cell Impedance)</small>	4.77	10 ¹² /L	3.8-4.8
 MCV <small>(Method: Calculated)</small>	87	fl	81-101
 MCH <small>(Method: Calculated)</small>	27.6	pg	27-32
 MCHC <small>(Method: Calculated)</small>	32.5	g/dL	32.5-34.5
 RDW-CV <small>(Method: Calculated)</small>	13.5	%	11.6-14.0
 Platelet Count (PLT) <small>(Method: Cell Impedance)</small>	251	10 ⁹ /L	150-410
 Total WBC Count <small>(Method: Impedance)</small>	9.2	10 ⁹ /L	4.0-10.0

Differential Leucocyte Count (DC)

 Neutrophils <small>(Method: Cell Impedance)</small>	66	%	40-70
 Lymphocytes <small>(Method: Cell Impedance)</small>	30	%	20-40
 Monocytes <small>(Method: Microscopy)</small>	02	%	2-10
 Eosinophils <small>(Method: Microscopy)</small>	02	%	1-6
 Basophils <small>(Method: Microscopy)</small>	00	%	1-2
 Absolute Neutrophils Count <small>(Method: Impedance)</small>	6.07	10 ⁹ /L	2.0-7.0
 Absolute Lymphocyte Count <small>(Method: Impedance)</small>	2.76	10 ⁹ /L	1.0-6.2
 Absolute Monocyte Count <small>(Method: Calculated)</small>	0.18	10 ⁹ /L	0.2-1.0
 Absolute Eosinophils Count <small>(Method: Calculated)</small>	0.18	10 ⁹ /L	0.02-0.5
 Absolute Basophil ICount <small>(Method: Calculated)</small>	0.00	10 ⁹ /L	0.0-0.3

Morphology
(Method: PAPS Staining)

Normocytic Normochromic



LABORATORY TEST REPORT

Name	: Miss. POOJA		
Sample ID	: A1309188		
Age/Gender	: 17 Years/Female	Reg. No	: 0312501160078
Referred by	: Dr. Nivedita Ashrit MD (Obs/Gyn)	SPP Code	: SPL-CV-172
Referring Customer	: V CARE MEDICAL DIAGNOSTICS	Collected On	: 16-Jan-2025 08:08 PM
Primary Sample	: Whole Blood	Received On	: 16-Jan-2025 10:50 PM
Sample Tested In	: Plasma-NaF(R)	Reported On	: 17-Jan-2025 12:41 AM
Client Address	: Kimtee colony ,Gokul Nagar,Tarnaka	Report Status	: Final Report


CLINICAL BIOCHEMISTRY
GLUCOSE RANDOM (RBS)

Test Name	Results	Units	Biological Reference Interval
-----------	---------	-------	-------------------------------

Glucose Random (RBS)	94	mg/dL	70-140
----------------------	----	-------	--------

(Method: Hexokinase (HK))

Interpretation of Plasma Glucose based on ADA guidelines 2018

Diagnosis	FastingPlasma Glucose(mg/dL)	2hrsPlasma Glucose(mg/dL)	HbA1c(%)	RBS(mg/dL)
Prediabetes	100-125	140-199	5.7-6.4	NA
Diabetes	> = 126	> = 200	> = 6.5	>=200(with symptoms)


Reference: Diabetes care 2018:41(suppl.1):S13-S27

- The random blood glucose if it is above 200 mg/dL and the patient has increased thirst, polyuria, and polyphagia, suggests diabetes mellitus.
- As a rule, two-hour glucose samples will reach the fasting level or it will be in the normal range.

*** End Of Report ***

Excellence In Health Care




DR.VAISHNAVI
MD BIOCHEMISTRY

Page 2 of 4

LABORATORY TEST REPORT

Name	: Miss. POOJA		
Sample ID	: A1309185, A1309186		
Age/Gender	: 17 Years/Female	Reg. No	: 0312501160078
Referred by	: Dr. Nivedita Ashrit MD (Obs/Gyn)	SPP Code	: SPL-CV-172
Referring Customer	: V CARE MEDICAL DIAGNOSTICS	Collected On	: 16-Jan-2025 08:08 PM
Primary Sample	: Whole Blood	Received On	: 16-Jan-2025 10:50 PM
Sample Tested In	: Whole Blood EDTA, Serum	Reported On	: 17-Jan-2025 12:29 AM
Client Address	: Kimtee colony ,Gokul Nagar,Tarnaka	Report Status	: Final Report



CLINICAL BIOCHEMISTRY

Test Name	Results	Units	Biological Reference Interval
Glycated Hemoglobin (HbA1c) <small>(Method: HPLC)</small>	5.0	%	Non Diabetic:< 5.7 Pre diabetic: 5.7-6.4 Diabetic:>= 6.5
Mean Plasma Glucose <small>(Method: Calculated)</small>	96.8	mg/dL	

Glycated hemoglobins (GHb), also called glycohemoglobins, are substances formed when glucose binds to hemoglobin, and occur in amounts proportional to the concentration of serum glucose. Since red blood cells survive an average of 120 days, the measurement of GHb provides an index of a person's average blood glucose concentration (glycemia) during the preceding 2-3 months. Normally, only 4% to 6% of hemoglobin is bound to glucose, while elevated glycohemoglobin levels are seen in diabetes and other hyperglycemic states Mean Plasma Glucose(MPG):This Is Mathematical Calculations Where Glycated Hb Can Be Correlated With Daily Mean Plasma Glucose Level

NOTE: The above Given Risk Level Interpretation is not age specific and is an information resource only and is not to be used or relied on for any diagnostic or treatment purposes and should not be used as a substitute for professional diagnosis and treatment. Kindly Correlate clinically.

INTERPRETATION

Method: Analyzer Fully automated HPLC platform.

Average Blood Glucose(eAG) (mg/dL)	Level of Control	Hemoglobin A1c (%)
421		14%
386		13%
350		12%
314		11%
279		10%
243		9%
208		8%
172	POOR	7%
136	GOOD	6%
101	EXCELLENT	5%

HbA1c values of 5.0- 6.5 percent indicate good control or an increased risk for developing diabetes mellitus. HbA1c values greater than 6.5 percent are diagnostic of diabetes mellitus. Diagnosis should be confirmed by repeating the HbA1c test.

NOTE: Hb F higher than 10 percent of total Hb may yield falsely low results. Conditions that shorten red cell survival, such as the presence of unstable hemoglobins like Hb SS, Hb CC, and Hb SC, or other causes of hemolytic anemia may yield falsely low results. Iron deficiency anemia may yield falsely high results.



Dr. Vaishnavi
DR.VAISHNAVI
MD BIOCHEMISTRY

LABORATORY TEST REPORT

Name	: Miss. POOJA		
Sample ID	: A1309185, A1309186		
Age/Gender	: 17 Years/Female	Reg. No	: 0312501160078
Referred by	: Dr. Nivedita Ashrit MD (Obs/Gyn)	SPP Code	: SPL-CV-172
Referring Customer	: V CARE MEDICAL DIAGNOSTICS	Collected On	: 16-Jan-2025 08:08 PM
Primary Sample	: Whole Blood	Received On	: 16-Jan-2025 10:50 PM
Sample Tested In	: Whole Blood EDTA, Serum	Reported On	: 17-Jan-2025 12:29 AM
Client Address	: Kimtee colony ,Gokul Nagar,Tarnaka	Report Status	: Final Report



CLINICAL BIOCHEMISTRY

Test Name	Results	Units	Biological Reference Interval
TSH -Thyroid Stimulating Hormone (Method: CLIA)	1.28	μIU/mL	0.35-5.5

Pregnancy & Cord Blood

TSH (Thyroid Stimulating Hormone (μIU/mL))	
First Trimester	: 0.24-2.99
Second Trimester	: 0.46-2.95
Third Trimester	: 0.43-2.78
Cord Blood	: 2.3-13.2

- TSH is synthesized and secreted by the anterior pituitary in response to a negative feedback mechanism involving concentrations of FT3 (free T3) and FT4 (free T4). Additionally, the hypothalamic tripeptide, thyrotropin-releasing hormone (TRH), directly stimulates TSH production.
- TSH interacts with specific cell receptors on the thyroid cell surface and exerts two main actions. The first action is to stimulate cell reproduction and hypertrophy. Secondly, TSH stimulates the thyroid gland to synthesize and secrete T3 and T4
- The ability to quantitate circulating levels of TSH is important in evaluating thyroid function. It is especially useful in the differential diagnosis of primary (thyroid) from secondary (pituitary) and tertiary (hypothalamus) hypothyroidism. In primary hypothyroidism, TSH levels are significantly elevated, while in secondary and tertiary hypothyroidism, TSH levels are low
- TRH stimulation differentiates secondary and tertiary hypothyroidism by observing the change in patient TSH levels. Typically, the TSH response to TRH stimulation is absent in cases of secondary hypothyroidism, and normal to exaggerated in tertiary hypothyroidism
- Historically, TRH stimulation has been used to confirm primary hyperthyroidism, indicated by elevated T3 and T4 levels and low or undetectable TSH levels. TSH assays with increased sensitivity and specificity provide a primary diagnostic tool to differentiate hyperthyroid from euthyroid patients.

*** End Of Report ***



Dr. Vaishnavi
DR. VAISHNAVI
MD BIOCHEMISTRY

Page 4 of 4