


LABORATORY TEST REPORT

Name	: Mrs. VENNELA		
Sample ID	: A1309341		
Age/Gender	: 26 Years/Female	Reg. No	: 0312501170041
Referred by	: Dr. Nivedita Ashrit MD (Obs/Gyn)	SPP Code	: SPL-CV-172
Referring Customer	: V CARE MEDICAL DIAGNOSTICS	Collected On	: 17-Jan-2025 06:18 PM
Primary Sample	: Whole Blood	Received On	: 17-Jan-2025 10:52 PM
Sample Tested In	: Serum	Reported On	: 18-Jan-2025 01:00 AM
Client Address	: Kimtee colony ,Gokul Nagar,Tarnaka	Report Status	: Final Report


CLINICAL BIOCHEMISTRY

Test Name	Results	Units	Biological Reference Interval
-----------	---------	-------	-------------------------------


Beta- Human Chorionic Gonadotropin Hormone <2.0 mIU/mL Refer to Interpretation
 (Method: CLIA)

Interpretation:

- A quantitative human chorionic gonadotropin (HCG) test measures the specific level of HCG in the blood. HCG is a hormone produced in the body during pregnancy.
- HCG appears in the blood and urine of pregnant women as early as 10 days after conception. Quantitative HCG measurement helps determine the exact age of the fetus. It can also assist in the diagnosis of abnormal pregnancies, such as ectopic pregnancies, molar pregnancies, and possible miscarriages. It is also used as part of a screening test for Down syndrome.
- This test is also done to diagnose abnormal conditions not related to pregnancy that can raise HCG level.


Non Pregnant Females: < 10.0 mIU/mL

Post Menopausal Females: < 10.0 mIU/mL

Pregnancy

Gestational Age and Expected hCG Values (mIU/mL)	Gestational Age and Expected hCG Values (mIU/mL)	Gestational Age and Expected hCG Values (mIU/mL)
0.2-1 weeks: 10-50	1-2 weeks : 50-500	2-3 weeks : 500-10,000
3-4 weeks : 1000-50,000	5-6 weeks : 10,000-100,000	6-8 weeks : 15,000-200,000
2-3 months : 10,000-100,000		




DR. VAISHNAVI
MD BIOCHEMISTRY

Page 1 of 3

LABORATORY TEST REPORT

Name	: Mrs. VENNELA		
Sample ID	: A1309341		
Age/Gender	: 26 Years/Female	Reg. No	: 0312501170041
Referred by	: Dr. Nivedita Ashrit MD (Obs/Gyn)	SPP Code	: SPL-CV-172
Referring Customer	: V CARE MEDICAL DIAGNOSTICS	Collected On	: 17-Jan-2025 06:18 PM
Primary Sample	: Whole Blood	Received On	: 17-Jan-2025 10:52 PM
Sample Tested In	: Serum	Reported On	: 18-Jan-2025 01:00 AM
Client Address	: Kimtee colony ,Gokul Nagar,Tarnaka	Report Status	: Final Report


CLINICAL BIOCHEMISTRY

Test Name	Results	Units	Biological Reference Interval
25 - Hydroxy Vitamin D <small>(Method: CLIA)</small>	9.42	ng/mL	<20.0-Deficiency 20.0-30.0-Insufficiency 30.0-100.0-Sufficiency >100.0-Potential Intoxication

Interpretation:

- Vitamin D helps your body absorb calcium and maintain strong bones throughout your entire life. Your body produces vitamin D when the sun's UV rays contact your skin. Other good sources of the vitamin include fish, eggs, and fortified dairy products. It's also available as a dietary supplement.
- Vitamin D must go through several processes in your body before your body can use it. The first transformation occurs in the liver. Here, your body converts vitamin D to a chemical known as 25-hydroxyvitamin D, also called calcidiol.
- The 25-hydroxy vitamin D test is the best way to monitor vitamin D levels. The amount of 25-hydroxyvitamin D in your blood is a good indication of how much vitamin D your body has. The test can determine if your vitamin D levels are too high or too low.
- The test is also known as the 25-OH vitamin D test and the calcidiol 25-hydroxycholecalciferol test. It can be an important indicator of osteoporosis (bone weakness) and rickets (bone malformation).

Those who are at high risk of having low levels of vitamin D include:

- people who don't get much exposure to the sun
- older adults
- people with obesity.
- dietary deficiency

Increased Levels: Vitamin D Intoxication

Method : CLIA

PRL(Prolactin) <small>(Method: CLIA)</small>	17.99	ng/mL	Refer Table
-------------------------------------------------	-------	-------	-------------

Interpretation:

Age	Reference Range: Male (ng/mL)	Reference Range: Female(ng/mL)
Puberty Tanner Stage		
1	< 10.0	3.6-12.0
2-3	< 6.1	2.6-18.0
4-5	2.8-11.0	3.2-20.0
Adult	2.1-17.7	Nonpregnant :2.8-29.2 Pregnant :9.7-208.5 Postmenopausal :1.8-20.3

- Prolactin is a 23kD sized hormone produced by the lactotroph cells of the pituitary gland, a grape-sized organ found at the base of the brain. Normally present in low amounts in men and non-pregnant women, prolactin's main role is to promote lactation (breast milk production).
- Breast milk production that is not related to childbirth (galactorrhea)
- Erection problems in men
- Irregular or no menstrual periods (amenorrhea)



Dr. Vaishnavi
DR. VAISHNAVI
MD BIOCHEMISTRY

Page 2 of 3

LABORATORY TEST REPORT

Name	: Mrs. VENNELA		
Sample ID	: A1309341		
Age/Gender	: 26 Years/Female	Reg. No	: 0312501170041
Referred by	: Dr. Nivedita Ashrit MD (Obs/Gyn)	SPP Code	: SPL-CV-172
Referring Customer	: V CARE MEDICAL DIAGNOSTICS	Collected On	: 17-Jan-2025 06:18 PM
Primary Sample	: Whole Blood	Received On	: 17-Jan-2025 10:52 PM
Sample Tested In	: Serum	Reported On	: 18-Jan-2025 01:00 AM
Client Address	: Kimtee colony ,Gokul Nagar,Tarnaka	Report Status	: Final Report


CLINICAL BIOCHEMISTRY

Test Name	Results	Units	Biological Reference Interval
Vitamin- B12 (cyanocobalamin) (Method: CLIA)	291	pg/mL	200-911

Interpretation:

This test is most often done when other blood tests suggest a condition called megaloblastic anemia. Pernicious anemia is a form of megaloblastic anemia caused by poor vitamin B12 absorption. This can occur when the stomach makes less of the substance the body needs to properly absorb vitamin B12.

Causes of vitamin B12 deficiency include: Diseases that cause malabsorption

- Lack of intrinsic factor, a protein that helps the intestine absorb vitamin B12
- Above normal heat production (for example, with hyperthyroidism)

An increased vitamin B12 level is uncommon in:

- Liver disease (such as cirrhosis or hepatitis)
- Myeloproliferative disorders (for example, polycythemia vera and chronic myelogenous leukemia)
-

 TSH -Thyroid Stimulating Hormone (Method: CLIA)	12.11	μIU/mL	0.35-5.5
---------------------------------------------------------------------------------------------------------------------------------------------	--------------	---------------	-----------------

Pregnancy & Cord Blood

TSH (Thyroid Stimulating Hormone (μIU/mL))	
First Trimester	: 0.24-2.99
Second Trimester	: 0.46-2.95
Third Trimester	: 0.43-2.78
Cord Blood	: 2.3-13.2

- TSH is synthesized and secreted by the anterior pituitary in response to a negative feedback mechanism involving concentrations of FT3 (free T3) and FT4 (free T4). Additionally, the hypothalamic tripeptide, thyrotropin-releasing hormone (TRH), directly stimulates TSH production.
- TSH interacts with specific cell receptors on the thyroid cell surface and exerts two main actions. The first action is to stimulate cell reproduction and hypertrophy. Secondly, TSH stimulates the thyroid gland to synthesize and secrete T3 and T4
- The ability to quantitate circulating levels of TSH is important in evaluating thyroid function. It is especially useful in the differential diagnosis of primary (thyroid) from secondary (pituitary) and tertiary (hypothalamus) hypothyroidism. In primary hypothyroidism, TSH levels are significantly elevated, while in secondary and tertiary hypothyroidism, TSH levels are low
- TRH stimulation differentiates secondary and tertiary hypothyroidism by observing the change in patient TSH levels. Typically, the TSH response to TRH stimulation is absent in cases of secondary hypothyroidism, and normal to exaggerated in tertiary hypothyroidism
- Historically, TRH stimulation has been used to confirm primary hyperthyroidism, indicated by elevated T3 and T4 levels and low or undetectable TSH levels. TSH assays with increased sensitivity and specificity provide a primary diagnostic tool to differentiate hyperthyroid from euthyroid patients.

*** End Of Report ***



Dr. Vaishnavi
DR. VAISHNAVI
MD BIOCHEMISTRY

Page 3 of 3