

LABORATORY TEST REPORT

Name	: Mrs. RASHMI		
Sample ID	: A1309986		
Age/Gender	: 39 Years/Female	Reg. No	: 0312502050013
Referred by	: Dr. SUNEETHA YERRAM	SPP Code	: SPL-CV-172
Referring Customer	: V CARE MEDICAL DIAGNOSTICS	Collected On	: 05-Feb-2025 10:21 AM
Primary Sample	: Whole Blood	Received On	: 05-Feb-2025 12:46 PM
Sample Tested In	: Plasma-NaF(R)	Reported On	: 05-Feb-2025 01:38 PM
Client Address	: Kimtee colony ,Gokul Nagar,Tarnaka	Report Status	: Final Report


CLINICAL BIOCHEMISTRY
GLUCOSE RANDOM (RBS)

Test Name	Results	Units	Biological Reference Interval
-----------	---------	-------	-------------------------------

Glucose Random (RBS)	104	mg/dL	70-140
----------------------	-----	-------	--------

(Method: Hexokinase (HK))

Interpretation of Plasma Glucose based on ADA guidelines 2018

Diagnosis	Fasting Plasma Glucose(mg/dL)	2hrs Plasma Glucose(mg/dL)	HbA1c(%)	RBS(mg/dL)
Prediabetes	100-125	140-199	5.7-6.4	NA
Diabetes	> = 126	> = 200	> = 6.5	>=200(with symptoms)

Reference: Diabetes care 2018;41(suppl.1):S13-S27

- The random blood glucose if it is above 200 mg/dL and the patient has increased thirst, polyuria, and polyphagia, suggests diabetes mellitus.
- As a rule, two-hour glucose samples will reach the fasting level or it will be in the normal range.

*** End Of Report ***

Excellence In Health Care



 DR. LAVANYA LAGISETTY
 MD BIOCHEMISTRY

LABORATORY TEST REPORT

Name	: Mrs. RASHMI		
Sample ID	: A1309985, A1309984		
Age/Gender	: 39 Years/Female	Reg. No	: 0312502050013
Referred by	: Dr. SUNEETHA YERRAM	SPP Code	: SPL-CV-172
Referring Customer	: V CARE MEDICAL DIAGNOSTICS	Collected On	: 05-Feb-2025 10:21 AM
Primary Sample	: Whole Blood	Received On	: 05-Feb-2025 12:24 PM
Sample Tested In	: Whole Blood EDTA, Serum	Reported On	: 05-Feb-2025 03:08 PM
Client Address	: Kimtee colony ,Gokul Nagar,Tarnaka	Report Status	: Final Report


CLINICAL BIOCHEMISTRY

Test Name	Results	Units	Biological Reference Interval
Glycated Hemoglobin (HbA1c) <small>(Method: HPLC)</small>	5.5	%	Non Diabetic:< 5.7 Pre diabetic: 5.7-6.4 Diabetic:>= 6.5
Mean Plasma Glucose <small>(Method: Calculated)</small>	111.15	mg/dL	

Glycated hemoglobins (GHb), also called glycohemoglobins, are substances formed when glucose binds to hemoglobin, and occur in amounts proportional to the concentration of serum glucose. Since red blood cells survive an average of 120 days, the measurement of GHb provides an index of a person's average blood glucose concentration (glycemia) during the preceding 2-3 months. Normally, only 4% to 6% of hemoglobin is bound to glucose, while elevated glycohemoglobin levels are seen in diabetes and other hyperglycemic states Mean Plasma Glucose(MPG):This Is Mathematical Calculations Where Glycated Hb Can Be Correlated With Daily Mean Plasma Glucose Level

NOTE: The above Given Risk Level Interpretation is not age specific and is an information resource only and is not to be used or relied on for any diagnostic or treatment purposes and should not be used as a substitute for professional diagnosis and treatment. Kindly Correlate clinically.

INTERPRETATION

Method: Analyzer Fully automated HPLC platform.

Average Blood Glucose(eAG) (mg/dL)	Level of Control	Hemoglobin A1c (%)
421		14%
386		13%
350		12%
314		11%
279		10%
243		9%
208		8%
172	POOR	7%
136	GOOD	6%
101	EXCELLENT	5%

HbA1c values of 5.0- 6.5 percent indicate good control or an increased risk for developing diabetes mellitus. HbA1c values greater than 6.5 percent are diagnostic of diabetes mellitus. Diagnosis should be confirmed by repeating the HbA1c test.

NOTE: Hb F higher than 10 percent of total Hb may yield falsely low results. Conditions that shorten red cell survival, such as the presence of unstable hemoglobins like Hb SS, Hb CC, and Hb SC, or other causes of hemolytic anemia may yield falsely low results. Iron deficiency anemia may yield falsely high results.



[Signature]
 DR. LAVANYA LAGISETTY
 MD BIOCHEMISTRY


LABORATORY TEST REPORT

Name	: Mrs. RASHMI		
Sample ID	: A1309985, A1309984		
Age/Gender	: 39 Years/Female	Reg. No	: 0312502050013
Referred by	: Dr. SUNEETHA YERRAM	SPP Code	: SPL-CV-172
Referring Customer	: V CARE MEDICAL DIAGNOSTICS	Collected On	: 05-Feb-2025 10:21 AM
Primary Sample	: Whole Blood	Received On	: 05-Feb-2025 12:24 PM
Sample Tested In	: Whole Blood EDTA, Serum	Reported On	: 05-Feb-2025 03:08 PM
Client Address	: Kimtee colony , Gokul Nagar, Tarnaka	Report Status	: Final Report




CLINICAL BIOCHEMISTRY

Test Name	Results	Units	Biological Reference Interval
-----------	---------	-------	-------------------------------

 **CA125 - Cancer Marker** 10.4 U/mL < 35.0
(Method: CLIA)

Interpretation:

The CA-125 blood test measures the level of the protein CA-125 in the blood. CA-125 is a protein that is found more in ovarian cancer cells than in other cells. This blood test is often used to monitor women who have been diagnosed with ovarian cancer. The test is useful if the CA-125 level was high when the cancer was first diagnosed. In these cases, measuring the CA-125 over time is a good tool to determine if ovarian cancer treatment is working. The CA-125 test may also be done if a woman has symptoms or findings on ultrasound that suggest ovarian cancer. In general, this test is not used to screen healthy women for ovarian cancer when a diagnosis has not yet been made. In a woman who has ovarian cancer, a rise in CA-125 usually means that the disease has progressed or come back (recurred). A decrease in CA-125 usually means the disease is responding to current treatment. In a woman who has not been diagnosed with ovarian cancer, a rise in CA-125 may mean a number of things. While it may mean that she has ovarian cancer, it can also indicate other types of cancer, as well as several other diseases, such as endometriosis, which are not cancer. In healthy women, an elevated CA-125 usually does not mean ovarian cancer is present. Most healthy women with an elevated CA-125 do not have ovarian cancer, or any other cancer. Any woman with an abnormal CA-125 test needs further tests. Sometimes surgery is needed to confirm the cause.

 **Carcino Embryonic Antigen (CEA)** 0.59 ng/mL Non Smokers:<3.00
(Method: CLIA) Smokers:<5.00

Interpretation:


- The carcinoembryonic antigen (CEA) test measures the level of CEA in the blood. CEA is a protein normally found in the tissue of a developing baby in the womb. The blood level of this protein disappears or becomes very low after birth. In adults, an abnormal level of CEA may be a sign of cancer.
- A high CEA level in a person recently treated for certain cancers may mean the cancer has returned.

A higher than normal level may be due to the following cancers:

Breast cancer
Cancers of the reproductive and urinary tracts
Colon cancer

An increased CEA level may also be due to:

Liver and gallbladder problems, such as scarring of the liver (cirrhosis), or gallbladder inflammation (cholecystitis)
Heavy smoking
Inflammatory bowel diseases (such as ulcerative colitis or diverticulitis)
Lung infection

 **TSH -Thyroid Stimulating Hormone** 2.45 µIU/mL 0.35-5.5
(Method: CLIA)

Pregnancy & Cord Blood

TSH (Thyroid Stimulating Hormone (µIU/mL))	
First Trimester	: 0.24-2.99
Second Trimester	: 0.46-2.95
Third Trimester	: 0.43-2.78
Cord Blood	: 2.3-13.2

- TSH is synthesized and secreted by the anterior pituitary in response to a negative feedback mechanism involving concentrations of FT3 (free T3) and FT4 (free T4). Additionally, the hypothalamic tripeptide, thyrotropin-releasing hormone (TRH), directly stimulates TSH production.
- TSH interacts with specific cell receptors on the thyroid cell surface and exerts two main actions. The first action is to stimulate cell reproduction and hypertrophy. Secondly, TSH stimulates the thyroid gland to synthesize and secrete T3 and T4
- The ability to quantitate circulating levels of TSH is important in evaluating thyroid function. It is especially useful in the differential diagnosis of primary (thyroid) from secondary (pituitary) and tertiary (hypothalamus) hypothyroidism. In primary hypothyroidism, TSH levels are significantly elevated, while in secondary and tertiary hypothyroidism, TSH levels are low
- TRH stimulation differentiates secondary and tertiary hypothyroidism by observing the change in patient TSH levels. Typically, the TSH response to TRH stimulation is absent in cases of secondary hypothyroidism, and normal to exaggerated in tertiary hypothyroidism
- Historically, TRH stimulation has been used to confirm primary hyperthyroidism, indicated by elevated T3 and T4 levels and low or undetectable TSH levels. TSH assays with increased sensitivity and specificity provide a primary diagnostic tool to differentiate hyperthyroid from euthyroid patients.




DR. LAVANYA LAGISSETTY
MD BIOCHEMISTRY

LABORATORY TEST REPORT

Name	: Mrs. RASHMI		
Sample ID	: A1309985, A1309984		
Age/Gender	: 39 Years/Female	Reg. No	: 0312502050013
Referred by	: Dr. SUNEETHA YERRAM	SPP Code	: SPL-CV-172
Referring Customer	: V CARE MEDICAL DIAGNOSTICS	Collected On	: 05-Feb-2025 10:21 AM
Primary Sample	: Whole Blood	Received On	: 05-Feb-2025 12:24 PM
Sample Tested In	: Whole Blood EDTA, Serum	Reported On	: 05-Feb-2025 03:08 PM
Client Address	: Kimtee colony ,Gokul Nagar,Tarnaka	Report Status	: Final Report



CLINICAL BIOCHEMISTRY

Test Name	Results	Units	Biological Reference Interval
-----------	---------	-------	-------------------------------

*** End Of Report ***




DR. LAVANYA LAGISETTY
MD BIOCHEMISTRY