

LABORATORY TEST REPORT

Name	: Mrs. V CHANDRAKALA		
Sample ID	: A0933983		
Age/Gender	: 67 Years/Female	Reg. No	: 0312502040038
Referred by	: Dr. SELF	SPP Code	: SPL-CV-172
Referring Customer	: V CARE MEDICAL DIAGNOSTICS	Collected On	: 04-Feb-2025 12:50 PM
Primary Sample	:	Received On	: 04-Feb-2025 04:00 PM
Sample Tested In	: Urine	Reported On	: 05-Feb-2025 08:37 AM
Client Address	: Kimtee colony ,Gokul Nagar,Tarnaka	Report Status	: Final Report


CLINICAL BIOCHEMISTRY

Test Name	Results	Units	Biological Reference Interval
Protein - Random Urine <small>(Method: Pyrogallol Red)</small>	112.5	mg/dL	1-14
Creatinine - Random Urine <small>(Method: kinetic Jaffe reaction.)</small>	460.98	mg/dL	15-278
Protein/Creatinine Ratio <small>(Method: Calculated)</small>	0.24		< 0.20

Interpretation:

The urine protein test measures the amount of protein being excreted in the urine. Proteinuria is frequently seen in chronic diseases, such as diabetes and hypertension, with increasing amounts of protein in the urine reflecting increasing kidney damage. With early kidney damage, the affected person is often asymptomatic. As damage progresses, or if protein loss is severe, the person may develop symptoms such as edema, shortness of breath, nausea, and fatigue. Excess protein overproduction, as seen with multiple myeloma, lymphoma, and amyloidosis, can also lead to proteinuria. Creatinine, a byproduct of muscle metabolism, is normally released into the urine at a constant rate.



 DR. LAVANYA LAGISETTY
 MD BIOCHEMISTRY

Page 1 of 2

LABORATORY TEST REPORT

Name	: Mrs. V CHANDRAKALA		
Sample ID	: A0933983		
Age/Gender	: 67 Years/Female	Reg. No	: 0312502040038
Referred by	: Dr. SELF	SPP Code	: SPL-CV-172
Referring Customer	: V CARE MEDICAL DIAGNOSTICS	Collected On	: 04-Feb-2025 12:50 PM
Primary Sample	:	Received On	: 04-Feb-2025 04:00 PM
Sample Tested In	: Urine	Reported On	: 06-Feb-2025 11:25 AM
Client Address	: Kimtee colony ,Gokul Nagar,Tarnaka	Report Status	: Final Report



MICROBIOLOGY

Culture and Sensitivity, Urine

Culture Comment No bacterial growth seen at the end of 48 hours of aerobic incubation.

ABST As per CLSI Guidelines.

Method : Aerobic Culture ABST; Disc Diffusion Method

*** End Of Report ***



Page 2 of 2



DR. RUTURAJ MANIKLAL KOLHAPURE
MD, MICROBIOLOGIST