

**LABORATORY TEST REPORT**

Name	: Mr. GOPI		
Sample ID	: A1840756		
Age/Gender	: 49 Years/Male	Reg. No	: 0312502130045
Referred by	: Dr. SELF	SPP Code	: SPL-CV-172
Referring Customer	: V CARE MEDICAL DIAGNOSTICS	Collected On	: 13-Feb-2025 05:42 PM
Primary Sample	: Whole Blood	Received On	: 13-Feb-2025 10:43 PM
Sample Tested In	: Serum	Reported On	: 14-Feb-2025 12:16 AM
Client Address	: Kimtee colony ,Gokul Nagar,Tarnaka	Report Status	: Final Report


**CLINICAL BIOCHEMISTRY**

Test Name	Results	Units	Biological Reference Interval
-----------	---------	-------	-------------------------------

Testosterone Total 214.24 ng/dL Refer Table  
 (Method: CLIA)

**Interpretation:**
**(Testosterone Reference Ranges)**

Age	Reference Range Male(ng/dL)	Reference Range Female(ng/dL)
Newborn(1-15days)	75-400	20-64
1-5 Months	1-177	1-5
6-11 Months	2-7	2-5
<b>Children:</b>		
1-5 Year	2-25	2-10
6-9 Year	3-30	2-20
<b>Puberty Tanner Stage</b>		
1	2-23	2-10
2	5-70	5-30
3	15-280	10-30
4	105-545	15-40
5	265-800	10-40
Adult	241-827	14-76

- Testosterone is a steroid hormone (androgen) made by the testes in males. Its production is stimulated and controlled by luteinising hormone (LH), which is manufactured in the pituitary gland. In males, testosterone stimulates development of secondary sex characteristics, including enlargement of the penis, growth of body hair and muscle, and a deepening voice. It is present in large amounts in males during puberty and in adult males to regulate the sex drive and maintain muscle mass. Testosterone is also produced by the adrenal glands in both males and females and, in small amounts, by the ovaries in females. The body can convert testosterone to oestradiol, the main sex hormone in females. There is great variability in testosterone levels between men and it is normal for testosterone levels to decline as men get older. Hypogonadism in a male refers to a reduction in sperm and/or testosterone production.



DR. LAVANYA LAGSETTY  
 MD BIOCHEMISTRY

**LABORATORY TEST REPORT**

Name	: Mr. GOPI		
Sample ID	: A1840756		
Age/Gender	: 49 Years/Male	Reg. No	: 0312502130045
Referred by	: Dr. SELF	SPP Code	: SPL-CV-172
Referring Customer	: V CARE MEDICAL DIAGNOSTICS	Collected On	: 13-Feb-2025 05:42 PM
Primary Sample	: Whole Blood	Received On	: 13-Feb-2025 10:43 PM
Sample Tested In	: Serum	Reported On	: 14-Feb-2025 12:16 AM
Client Address	: Kimtee colony ,Gokul Nagar,Tarnaka	Report Status	: Final Report


**CLINICAL BIOCHEMISTRY**

Test Name	Results	Units	Biological Reference Interval
FSH (Follicle Stimulating Hormone) <small>(Method: CLIA)</small>	6.66	mIU/mL	Refer Table

**Interpretation:**

Age	Reference Range: Male (mIU/mL)	Reference Range: Female(mIU/mL)
<b>Pre Puberty Child</b>		
2-11 Months	0.19-11.3	0.10-11.3
1-10 Years	0.3-4.6	0.68-6.7
<b>Puberty Tanner Stage</b>		
1-2	0.30-4.6	0.68-6.7
3-4	1.24-15.4	1.0-7.4
5	1.53-6.8	1.0-9.2
<b>Adult</b>	1.42-18.4	
Follicular Phase	----	2.5-10.2
Midcycle Peak	----	3.4-33.4
Luteal Phase	----	1.5-9.1
Postmenopausal	----	23.0-116.3
Pregnant	----	< 0.3

The follicle stimulating hormone (FSH) blood test measures the level of FSH in blood. FSH is a hormone released by the pituitary gland, located on the underside of the brain.

**Low FSH levels in women may be present due to:**

- Being very underweight or having had recent rapid weight loss
- Not producing eggs (not ovulating)
- Parts of the brain (the pituitary gland or hypothalamus) not producing normal amounts of some or all of its hormones
- Pregnancy

**High FSH levels in men may mean the testicles are not functioning correctly due to:**

- Advancing age (male menopause)
- Damage to testicles caused by alcohol abuse, chemotherapy, or radiation
- Certain tumors in the pituitary gland

\*\*\* End Of Report \*\*\*



*[Signature]*  
 DR. LAVANYA LAGISETTY  
 MD BIOCHEMISTRY

Page 2 of 2