










**LABORATORY TEST REPORT**

|                    |                                      |               |                        |
|--------------------|--------------------------------------|---------------|------------------------|
| Name               | : Mrs. SOUNDARIYA                    |               |                        |
| Sample ID          | : A1840801                           |               |                        |
| Age/Gender         | : 26 Years/Female                    | Reg. No       | : 0312502140029        |
| Referred by        | : Dr. Nivedita Ashrit MD (Obs/Gyn)   | SPP Code      | : SPL-CV-172           |
| Referring Customer | : V CARE MEDICAL DIAGNOSTICS         | Collected On  | : 14-Feb-2025 12:19 PM |
| Primary Sample     | : Whole Blood                        | Received On   | : 14-Feb-2025 04:23 PM |
| Sample Tested In   | : Whole Blood EDTA                   | Reported On   | : 14-Feb-2025 04:36 PM |
| Client Address     | : Kimtee colony ,Gokul Nagar,Tarnaka | Report Status | : Final Report         |












**HAEMATOLOGY**

| Test Name | Results | Units | Biological Reference Interval |
|-----------|---------|-------|-------------------------------|
|-----------|---------|-------|-------------------------------|

**Complete Blood Picture(CBP)**

|   |             |                     |           |
|---|-------------|---------------------|-----------|
|  <b>Haemoglobin (Hb)</b><br>(Method: Cymeth Method)        | 13.2        | g/dL                | 12-15     |
|  <b>Haematocrit (HCT)</b><br>(Method: Calculated)          | <b>37.7</b> | %                   | 40-50     |
|  <b>RBC Count</b><br>(Method: Cell Impedance)              | <b>4.85</b> | 10 <sup>12</sup> /L | 3.8-4.8   |
|  <b>MCV</b><br>(Method: Calculated)                        | <b>78</b>   | fl                  | 81-101    |
|  <b>MCH</b><br>(Method: Calculated)                       | 27.3        | pg                  | 27-32     |
|  <b>MCHC</b><br>(Method: Calculated)                     | 33.0        | g/dL                | 32.5-34.5 |
|  <b>RDW-CV</b><br>(Method: Calculated)                   | <b>15.2</b> | %                   | 11.6-14.0 |
|  <b>Platelet Count (PLT)</b><br>(Method: Cell Impedance) | <b>412</b>  | 10 <sup>9</sup> /L  | 150-410   |
|  <b>Total WBC Count</b><br>(Method: Impedance)           | 7.8         | 10 <sup>9</sup> /L  | 4.0-10.0  |

**Differential Leucocyte Count (DC)**

|   |      |                    |          |
|---|------|--------------------|----------|
|  <b>Neutrophils</b><br>(Method: Cell Impedance)            | 65   | %                  | 40-70    |
|  <b>Lymphocytes</b><br>(Method: Cell Impedance)            | 25   | %                  | 20-40    |
|  <b>Monocytes</b><br>(Method: Microscopy)                  | 06   | %                  | 2-10     |
|  <b>Eosinophils</b><br>(Method: Microscopy)                | 04   | %                  | 1-6      |
|  <b>Basophils</b><br>(Method: Microscopy)                  | 00   | %                  | 1-2      |
|  <b>Absolute Neutrophils Count</b><br>(Method: Impedance)  | 5.07 | 10 <sup>9</sup> /L | 2.0-7.0  |
|  <b>Absolute Lymphocyte Count</b><br>(Method: Impedance)   | 1.95 | 10 <sup>9</sup> /L | 1.0-3.0  |
|  <b>Absolute Monocyte Count</b><br>(Method: Calculated)    | 0.47 | 10 <sup>9</sup> /L | 0.2-1.0  |
|  <b>Absolute Eosinophils Count</b><br>(Method: Calculated) | 0.31 | 10 <sup>9</sup> /L | 0.02-0.5 |
|  <b>Absolute Basophil ICount</b><br>(Method: Calculated)   | 0.00 | 10 <sup>9</sup> /L | 0.0-0.3  |

**Morphology**  
 (Method: PAPs Staining)

Anisocytosis with Normocytic normochromic with Mild Thrombocytosis



**LABORATORY TEST REPORT**

|                    |                                      |               |                        |
|--------------------|--------------------------------------|---------------|------------------------|
| Name               | : Mrs. SOUNDARIYA                    |               |                        |
| Sample ID          | : A1840800                           |               |                        |
| Age/Gender         | : 26 Years/Female                    | Reg. No       | : 0312502140029        |
| Referred by        | : Dr. Nivedita Ashrit MD (Obs/Gyn)   | SPP Code      | : SPL-CV-172           |
| Referring Customer | : V CARE MEDICAL DIAGNOSTICS         | Collected On  | : 14-Feb-2025 12:19 PM |
| Primary Sample     | : Whole Blood                        | Received On   | : 14-Feb-2025 04:23 PM |
| Sample Tested In   | : Serum                              | Reported On   | : 14-Feb-2025 06:33 PM |
| Client Address     | : Kimtee colony ,Gokul Nagar,Tarnaka | Report Status | : Final Report         |


**CLINICAL BIOCHEMISTRY**

| Test Name | Results | Units | Biological Reference Interval |
|-----------|---------|-------|-------------------------------|
|-----------|---------|-------|-------------------------------|


**TSH -Thyroid Stimulating Hormone**
0.84
μIU/mL
0.35-5.5

(Method: CLIA)

**Pregnancy & Cord Blood**

| TSH (Thyroid Stimulating Hormone (μIU/mL)) |             |
|--|-------------|
| First Trimester                            | : 0.24-2.99 |
| Second Trimester                           | : 0.46-2.95 |
| Third Trimester                            | : 0.43-2.78 |
| Cord Blood                                 | : 2.3-13.2  |

- TSH is synthesized and secreted by the anterior pituitary in response to a negative feedback mechanism involving concentrations of FT3 (free T3) and FT4 (free T4). Additionally, the hypothalamic tripeptide, thyrotropin-releasing hormone (TRH), directly stimulates TSH production.
- TSH interacts with specific cell receptors on the thyroid cell surface and exerts two main actions. The first action is to stimulate cell reproduction and hypertrophy. Secondly, TSH stimulates the thyroid gland to synthesize and secrete T3 and T4
- The ability to quantitate circulating levels of TSH is important in evaluating thyroid function. It is especially useful in the differential diagnosis of primary (thyroid) from secondary (pituitary) and tertiary (hypothalamus) hypothyroidism. In primary hypothyroidism, TSH levels are significantly elevated, while in secondary and tertiary hypothyroidism, TSH levels are low
- TRH stimulation differentiates secondary and tertiary hypothyroidism by observing the change in patient TSH levels. Typically, the TSH response to TRH stimulation is absent in cases of secondary hypothyroidism, and normal to exaggerated in tertiary hypothyroidism
- Historically, TRH stimulation has been used to confirm primary hyperthyroidism, indicated by elevated T3 and T4 levels and low or undetectable TSH levels. TSH assays with increased sensitivity and specificity provide a primary diagnostic tool to differentiate hyperthyroid from euthyroid patients.

\*\*\* End Of Report \*\*\*



  
 DR. LAVANYA LAGISETTY  
 MD BIOCHEMISTRY

Page 2 of 2