

Sagepath Labs Pvt. Ltd.

Lab Address: - # Plot No. 564, 1st floor, Buddhanagar, Near Sai Baba Temple Peerzadiguda Boduppal Hyderabad, Telangana. ICMR Reg .No. SAPALAPVLHT (Covid -19)

REPORT LABORATORY TEST

Name : Mrs. RADHIKA Sample ID : A1841627

> Reg. No : 0312503040022

Referred by : Dr. SELF SPP Code : SPL-CV-172

: 04-Mar-2025 02:09 PM

Collected On

Referring Customer: V CARE MEDICAL DIAGNOSTICS Primary Sample : Whole Blood

Received On : 04-Mar-2025 04:25 PM

Sample Tested In : Serum

: 28 Years/Female

Reported On : 04-Mar-2025 06:48 PM

Client Address : Kimtee colony ,Gokul Nagar,Tarnaka Report Status : Final Report

	IMMUNOLOGY & SEROLOGY		
Test Name	Results	Units	Biological Reference Interval
Hepatitis C Virus Antibody	0.17	S/Co	< 1.00 : Negative > 1.00 : Positive


Interpretation:

Age/Gender

- 1. Negative result implies that antibodies to HCV have not been detected in the sample. This means the patient has either not been exposed to HCV infection or the sample has been tested during the "window phase" i.e. before the development of detectable levels of antibodies. Hence a Non-Reactive result does not exclude the possibility of exposure or infection with HCV.
- 2. Positive result implies that antibodies to HCV have been detected in the sample.

Comments :-

Hepatitis C (HCV) is an RNA virus of Flavivirus group transmitted via blood transfusions, transplantation, injection drug users, accidental needle punctures in healthcare workers, dialysis patients and rarely from mother to infant. 10% of new cases show sexual transmission. As compared to HAV & HBV, chronic infection with HCV occurs in 85% of infected individuals. In high risk populations, the predictive value of Anti HCV for HCV infection is > 99% whereas in low risk populations it is only 25%



- 1. False positive results are seen in Autoimmune diseases, Rheumatoid factor, Hypergammaglobulinemia, Paraproteinemia, passive antibody transfer, Anti- idiotypes
- 2. False negative results are seen in early Acute infection, Immunosuppression & Immuno-incompetence
- 3. HCV RNA PCR recommended in all Reactive results to differentiate between past and present infection

*** End Of Report ***







DR. RUTURAJ MANIKLAL KOLHAPURE MD, MICROBIOLOGIST