Sample ID : A1841791

Age/Gender : 60 Years/Female Reg. No : 0312503100002

Referred by : Dr. PRADEEP KUMAR MISHRA SPP Code : SPL-CV-172

Referring Customer : V CARE MEDICAL DIAGNOSTICS Collected On : 10-Mar-2025 08:46 AM
Primary Sample : Whole Blood Received On : 10-Mar-2025 01:06 PM
Sample Tested In : Whole Blood EDTA Reported On : 10-Mar-2025 01:33 PM

Client Address : Kimtee colony , Gokul Nagar, Tarnaka Report Status : Final Report

HAEMATOLOGY SAGE CARE 1.2 Test Name Results Units Biological Reference Interval

Test Name	Results	Units	Biological Reference Interval
COMPLETE BLOOD COUNT (CBC)			
Haemoglobin (Hb) (Method: Cymreth Method)	<u>11.6</u>	g/dL	12-15
(Wellock: Cyriment method) (Refloc Count (Method: Cell Impedence)	3.86	10^12/L	3.8-4.8
Haematocrit (HCT)	<u>35.6</u>	%	40-50
(Method: Calculated) MCV	92	fl	81-101
(Method: Calculated) MCH	30.0	pg	27-32
(Method: Calculated) MCHC (Method: Calculated)	32.6	g/dL	32.5-34.5
RDW-CV (Method: Calculated)	13.1	%	11.6-14.0
(Method: Call Impedance) (Method: Cell Impedance)	290	10^9/L	150-410
(Method: Cell Impedance) Total WBC Count (Method: Impedance)	6.3	10^9/L	4.0-10.0
(Method: Impedance) Neutrophils (Method: Cell Impedence)	61	%	40-70
Absolute Neutrophils Count	3.84	10^9/L	2.0-7.0
(Method: Impedence) Lymphocytes (Method: Cell Impedence)	32	%	20-40
Absolute Lymphocyte Count (Method: Impedence)	2.02	10^9/L	1.0-3.0
(Method: Impedence) Monocytes (Method: Microscopy)	05	%	2-10
(Method: Macroscopy) Absolute Monocyte Count (Method: Calculated)	0.32	10^9/L	0.2-1.0
Cosinophils (Method: Microscopy)	02	%	1-6
(Method: Microscopy) Absolute Eosinophils Count (Method: Calculated)	0.13	10^9/L	0.02-0.5
Basophils	00	%	1-2
(Method: Microscopy) Absolute Basophil ICount	0.00	10^9/L	0.0-0.3
(Method: Calculated) Atypical cells	0.00		
<u>Morphology</u>			
WBC	Within Nor	mal Limits	
RBC	Normocytic	c normochromic	c blood picture.
Platelets	Adequate.		

Platelets Adequate.







Sample ID : A1841791

Age/Gender : 60 Years/Female Reg. No : 0312503100002

Referred by : Dr. PRADEEP KUMAR MISHRA SPP Code : SPL-CV-172

Referring Customer : V CARE MEDICAL DIAGNOSTICS Collected On : 10-Mar-2025 08:46 AM
Primary Sample : Whole Blood Received On : 10-Mar-2025 01:06 PM
Sample Tested In : Whole Blood EDTA Reported On : 10-Mar-2025 02:14 PM

Client Address : Kimtee colony ,Gokul Nagar,Tarnaka Report Status : Final Report

HAEMATOLOGY

SAGE CARE 1.2

Test Name	Results	Units	Biological Reference Interval

Erythrocyte Sedimentation Rate (ESR)
15
mm/hr 12 or less

Comments: ESR is an acute phase reactant which indicates presence and intensity of an inflammatory process. It is never diagnostic of a specific disease. It is used to monitor the course or response to treatment of certain diseases. Extremely high levels are found in cases of malignancy, hematologic diseases, collagen disorders and renal diseases.







Sample ID : A1841790

Age/Gender : 60 Years/Female Reg. No : 0312503100002

Referred by : Dr. PRADEEP KUMAR MISHRA SPP Code : SPL-CV-172

Referring Customer : V CARE MEDICAL DIAGNOSTICS Collected On : 10-Mar-2025 08:46 AM Primary Sample Received On : 10-Mar-2025 03:45 PM

Sample Tested In : Urine Reported On : 10-Mar-2025 04:35 PM Client Address : Kimtee colony ,Gokul Nagar,Tarnaka Report Status : Final Report

(•	LI	N	IJ	С	Α	L	Р	Α	Т	Н	Ω	L	O	G'	Υ
				٠.	•		_		•					•	_	

Test Name	Results	Units	Biological Reference Interval

Complete Urine Analysis (CUE)

Physical Examination

Colour Pale Yellow Straw to light amber

HAZY Clear **Appearance**

Chemical Examination Glucose Negative Negative Protein Negative Negative Negative Negative Bilirubin (Bile) Negative Negative Urobilinogen Ketone Bodies Negative Negative Specific Gravity 1.020 1.000 - 1.030 Blood Negative Negative 6.0 5.0 - 8.5 Reaction (pH) Negative **Nitrites** Negative Leukocyte esterase (+) Negative Microscopic Examination (Microscopy) PUS(WBC) Cells 06-08 /hpf 00-05 R.B.C. Nil /hpf Nil **Epithelial Cells** 02-03 /hpf 00-05 Absent Absent Casts

Absent

Nil

Nil

Comments: Urine analysis is one of the most useful laboratory tests as it identifies a wide range of medical conditions including renal damage, urinary tract infections, diabetes, hypertension and drug toxicity.

Absent

Absent

Nil



Crystals Bacteria

Budding Yeast Cells





Name : Mrs. S NAGAMANI Sample ID : A1841794, A1841789

No : 0312503100002

Age/Gender : 60 Years/Female Reg. No : 0312503100002 Referred by : Dr. PRADEEP KUMAR MISHRA SPP Code : SPL-CV-172

Referring Customer : V CARE MEDICAL DIAGNOSTICS Collected On : 10-Mar-2025 08:46 AM
Primary Sample : Whole Blood Received On : 10-Mar-2025 03:45 PM
Sample Tested In : Plasma-NaF(F), Plasma-NaF(PP) Reported On : 10-Mar-2025 04:32 PM

Client Address : Kimtee colony ,Gokul Nagar,Tarnaka Report Status : Final Report

CLINICAL BIOCHEMISTRY

GLUCOSE POST PRANDIAL (PP)

Test Name Results Units Biological Reference Interval

Glucose Fasting (F) <u>122</u> mg/dL 70-100

Interpretation of Plasma Glucose based on ADA guidelines 2018

Diagnosis	FastingPlasma Glucose(mg/dL)	2hrsPlasma Glucose(mg/dL)	HbA1c(%)	RBS(mg/dL)
Prediabetes	100-125	140-199	5.7-6.4	NA
Diabetes	> = 126	>= 200	>= 6.5	>=200(with symptoms)

Reference: Diabetes care 2018:41(suppl.1):S13-S27

Glucose Post Prandial (PP) 120 mg/dL 70-140

(Method: Hexokinase (HK))

 Interpretation of Plasma Glucose based on ADA guidelines 2018

 Diagnosis
 FastingPlasma Glucose(mg/dL)
 2hrsPlasma Glucose(mg/dL)
 HbA1c(%)
 RBS(mg/dL)

 Prediabetes
 100-125
 140-199
 5.7-6.4
 NA

 Diabetes
 > = 126
 > = 200
 > = 6.5
 >=200(with symptoms)

Reference: Diabetes care 2018:41(suppl.1):S13-S27

- · Postprandial glucose level is a screening test for Diabetes Mellitus
- If glucose level is >140 mg/dL and <200 mg/dL, then GTT (glucose tolerance test) is advised.
- \bullet If level after 2 hours = >200 mg/dL diabetes mellitus is confirmed.
- Advise HbA1c for further evaluation.









Name : Mrs. S NAGAMANI

Sample ID : A1841791

Age/Gender : 60 Years/Female Reg. No : 0312503100002

Referred by : Dr. PRADEEP KUMAR MISHRA SPP Code : SPL-CV-172

Referring Customer : V CARE MEDICAL DIAGNOSTICS Collected On : 10-Mar-2025 08:46 AM
Primary Sample : Whole Blood Received On : 10-Mar-2025 01:06 PM
Sample Tested In : Whole Blood EDTA Reported On : 10-Mar-2025 02:08 PM

Client Address : Kimtee colony ,Gokul Nagar,Tarnaka Report Status : Final Report

CLINICAL BIOCHEMISTRY

SAGE CARE 1.2

SAGE CARE 1.2						
Test Name	Results	Units	Biological Reference Interval			
Glycated Hemoglobin (HbA1c)	<u>7.1</u>	%	Non Diabetic:< 5.7 Pre diabetic: 5.7-6.4 Diabetic:>= 6.5			
Mean Plasma Glucose	157.07	mg/dL				

Glycated hemoglobins (GHb), also called glycohemoglobins, are substances formed when glucose binds to hemoglobin, and occur in amounts proportional to the concentration of serum glucose. Since red blood cells survive an average of 120 days, the measurement of GHb provides an index of a person's average blood glucose concentration (glycemia) during the preceding 2-3 months. Normally, only 4% to 6% of hemoglobin is bound to glucose, while elevated glycohemoglobin levels are seen in diabetes and other hyperglycemic states Mean Plasma Glucose (MPG): This Is Mathematical Calculations Where Glycated Hb Can Be Correlated With Daily Mean Plasma Glucose Level

NOTE: The above Given Risk Level Interpretation is not age specific and is an information resource only and is not to be used or relied on for any diagnostic or treatment purposes and should not be used as a substitute for professional diagnosis and treatment. Kindly Correlate clinically.

INTERPRETATION

Method: Analyzer Fully automated HPLC platform.

Average Blood Glucose(eAG) (mg/dL)	Level of Control	Hemoglobin A1c (%)
421		14%
386	_ A	13%
350	L	12%
314	E	11%
279	R	10%
243	Т	9%
208		8%
172	POOR	7%
136	GOOD	6%
101	EXCELLENT	5%

HbA1c values of 5.0- 6.5 percent indicate good control or an increased risk for developing diabetes mellitus. HbA1c values greater than 6.5 percent are diagnostic of diabetes mellitus. Diagnosis should be confirmed by repeating the HbA1c test.

NOTE: Hb F higher than 10 percent of total Hb may yield falsely low results. Conditions that shorten red cell survival, such as the presence of unstable hemoglobins like Hb SS, Hb CC, and Hb SC, or other causes of hemolytic anemia may yield falsely low results. Iron deficiency anemia may yield falsely high results.









Sample ID : A1841792

Age/Gender : 60 Years/Female Reg. No : 0312503100002

Referred by : Dr. PRADEEP KUMAR MISHRA SPP Code : SPL-CV-172

Referring Customer : V CARE MEDICAL DIAGNOSTICS Collected On : 10-Mar-2025 08:46 AM
Primary Sample : Whole Blood Received On : 10-Mar-2025 01:06 PM
Sample Tested In : Serum Reported On : 10-Mar-2025 03:23 PM

Client Address : Kimtee colony ,Gokul Nagar,Tarnaka Report Status : Final Report

CLINICAL BIOCHEMISTRY SAGE CARE 1.2 Test Name Results Units Biological Reference Interval © Calcium (Method: Arsenazo) 9.1 mg/dL 8.5-10.1

Comments:

- Calcium in the body is found mainly in the bones (approximately 99%). In serum, Calcium exists in a
 free ionised form and in bound form (with Albumin). Hence, a decrease in Albumin causes lower
 Calcium levels and vice-versa.
- Calcium levels in serum depend on the Parathyroid Hormone.
- Increased Calcium levels are found in Bone tumors, Hyperparathyroidism. decreased levels are found in Hypoparathyroidism, renal failure, Rickets.









Referring Customer : V CARE MEDICAL DIAGNOSTICS

Primary Sample : Whole Blood Sample Tested In : Serum

Client Address : Kimtee colony ,Gokul Nagar,Tarnaka

Lab Address:- # Plot No. 564 , 1st floor , Buddhanagar , Near Sai Baba Temple Peerzadiguda Boduppal Hyderabad, Telangana.

: 0312503100002 : SPL-CV-172

Collected On : 10-Mar-2025 08:46 AM

Received On : 10-Mar-2025 01:06 PM

Reported On : 10-Mar-2025 03:23 PM

Report Status : Final Report

ICMR Reg .No. SAPALAPVLHT (Covid -19)

Reg. No

SPP Code

CLINICAL BIOCHEMISTRY

SAGE CARE 1.2

SAGE CARE 1.2							
Test Name	Results	Units	Biological Reference Interval				
Lipid Profile							
Cholesterol Total	171	mg/dL	< 200				
Triglycerides-TGL (Method: GPO-POD)	82	mg/dL	< 150				
© Cholesterol-HDL (Method: Direct)	42	mg/dL	40-60				
© Cholesterol-LDL (Method: Calculated)	<u>112.6</u>	mg/dL	< 100				
Cholesterol- VLDL (Method: Calculated)	16.4	mg/dL	7-35				
Non HDL Cholesterol (Method: Calculated)	129	mg/dL	< 130				
© Cholesterol Total /HDL Ratio	<u>4.07</u>	Ratio	0-4.0				
LDL/HDL Ratio (Method: Calculated)	2.68	Ratio	0-3.5				

The National Cholesterol Education program's third Adult Treatment Panel (ATPIII) has issued its recommendations on evaluating and treating lipid discorders for primary and secondary.

NCEP Recommendations	Cholesterol Total in (mg/dL)	Triglycerides	HDL Cholesterol (mg/dL)	LDL Cholesterol in (mg/dL)	Non HDL Cholesterol in (mg/dL)
Untimai	Adult: < 200 Children: < 170	< 150	40-59	Adult:<100 Children: <110	<130
Above Optimal				100-129	130 - 159
Borderline High	Adult: 200-239 Children:171-199	150-199		Adult: 130-159 Children: 111-129	160 - 189
High	Adult:>or=240 Children:>or=200	200-499	≥ 60	Adult:160-189 Children:>or=130	190 - 219
Very High		>or=500		Adult: >or=190	>=220

Note: LDL cholesterol cannot be calculated if triglyceride is >400 mg/dL (Friedewald's formula). Calculated values not provided for LDL and VLDL









: 0312503100002

Sample ID : A1841792

> : 60 Years/Female : Dr. PRADEEP KUMAR MISHRA

SPP Code : SPL-CV-172

ICMR Reg .No. SAPALAPVLHT (Covid -19)

Referred by

Referring Customer: V CARE MEDICAL DIAGNOSTICS

Collected On : 10-Mar-2025 08:46 AM Received On : 10-Mar-2025 01:06 PM

Lab Address: - # Plot No. 564, 1st floor, Buddhanagar, Near Sai Baba Temple Peerzadiguda Boduppal Hyderabad, Telangana.

Primary Sample : Whole Blood Sample Tested In : Serum

Reported On : 10-Mar-2025 03:23 PM

Client Address : Kimtee colony ,Gokul Nagar,Tarnaka

Report Status : Final Report

CLINICAL BIOCHEMISTRY

SAGE CARE 12

SAGE CARE 1.2								
Test Name	Results	Units	Biological Reference Interval					
Liver Function Test (LFT)								
Bilirubin(Total)	0.4	mg/dL	0.3-1.2					
Bilirubin (Direct)	0.1	mg/dL	0.0 - 0.3					
Bilirubin (Indirect)	0.3	mg/dL	0.2-1.0					
Aspartate Aminotransferase (AST/SGOT)	18	U/L	15-37					
Alanine Aminotransferase (ALT/SGPT)	10	U/L	0-55					
Malkaline Phosphatase(ALP)	60	U/L	30-120					
(GGTP)	10	U/L	5-55					
Method: Bluret)	7.1	g/dL	6.4-8.2					
Albumin (Method: Bromocresol Green (BCG))	4.0	g/dL	3.4-5.0					
(Matthew Sales Gent (p.C.)) (Matthew Sales Gent (p.C.))	3.1	g/dL	2.0-4.2					
A:G Ratio Method: Calculated)	1.29	Ratio	0.8-2.0					
SGOT/SGPT Ratio	<u>1.8</u>	Ratio	<1.0					

Alanine Aminotransferase(ALT) is an enzyme found in liver and kidneys cells. ALT helps create energy for liver cells. Damaged liver cells release ALT into the bloodstream, which can elevate ALT levels in the blood.

Aspartate Aminotransferase (AST) is an enzyme in the liver and muscles that helps metabolizes amino acids. Similarly to ALT, elevated AST levels may be a sign of liver damage or liver disease.

Alkaline phosphate (ALP) is an enzyme present in the blood. ALP contributes to numerous vital bodily functions, such as supplying nutrients to the liver, promoting bone growth, and metabolizing fat in the intestines.

Gamma-glutamyl Transpeptidase (GGTP) is an enzyme that occurs primarily in the liver, but it is also present in the kidneys, pancreas, gallbladder, and spleen. Higher than normal concentrations of GGTP in the blood may indicate alcohol-related liver damage. Elevated GGTP levels can also increase the risk of developing certain types of cancer.

Bilirubin is a waste product that forms when the liver breaks down red blood cells. Bilirubin exits the body as bile in stool. High levels of bilirubin can cause jaundice - a condition in which the skin and whites of the eyes turn yellow- and may indicate liver damage. Albumin is a protein that the liver produces. The liver releases albumin into the bloodstream, where it helps fight infections and transport vitamins, hormones, and enzymes throughout the body.

*** End Of Report ***





Liver damage can cause abnormally low albumin levels.





Sample ID : A1841792

Age/Gender : 60 Years/Female Reg. No : 0312503100002

Referred by : Dr. PRADEEP KUMAR MISHRA SPP Code : SPL-CV-172

Referring Customer : V CARE MEDICAL DIAGNOSTICS Collected On : 10-Mar-2025 08:46 AM
Primary Sample : Whole Blood Received On : 10-Mar-2025 01:06 PM
Sample Tested In : Serum Reported On : 10-Mar-2025 03:23 PM

Client Address : Kimtee colony ,Gokul Nagar,Tarnaka Report Status : Final Report

CLINICAL BIOCHEMISTRY

SAGE CARE 1

	SAGE CARE 1.2						
Test Name	Results	Units	Biological Reference Interval				
Kidney Profile-KFT							
Creatinine (Method: Sarcosine Oxidase Method)	0.78	mg/dL	0.60-1.10				
Urea-Serum (Method: Urease-GLDH, UV Method)	31.4	mg/dL	12.8-42.8				
Blood Urea Nitrogen (BUN)	14.67	mg/dL	7.0-18.0				
BUN / Creatinine Ratio	18.81	Ratio	6 - 22				
Uric Acid (Method: Uricase)	4.3	mg/dL	2.6-6.0				
Sodium (Method: 15E Direct)	143	mmol/L	135-150				
Potassium (Method: ISE Direct)	4.1	mmol/L	3.5-5.0				
Chloride Method: ISE Direct)	102	mmol/L	94-110				

Interpretation:

• The kidneys, located in the retroperitoneal space in the abdomen, are vital for patient health. They process several hundred liters of fluid a day and remove around two liters of waste products from the bloodstream. The volume of fluid that passes though the kidneys each minute is closely linked to cardiac output. The kidneys maintain the body's balance of water and concentration of minerals such as sodium, potassium, and phosphorus in blood and remove waste by-products from the blood after digestion, muscle activity and exposure to chemicals or medications. They also produce renin which helps regulate blood pressure, produce erythropoietin which stimulates red blood cell production, and produce an active form of vitamin D, needed for bone health.









Name : Mrs. S NAGAMANI

Sample ID : A1841792

Age/Gender : 60 Years/Female Reg. No : 0312503100002

Referred by : Dr. PRADEEP KUMAR MISHRA SPP Code : SPL-CV-172

Referring Customer : V CARE MEDICAL DIAGNOSTICS Collected On : 10-Mar-2025 08:46 AM

Primary Sample : Whole Blood Received On : 10-Mar-2025 01:06 PM

Sample Tested In : Serum Reported On : 10-Mar-2025 03:23 PM

Client Address : Kimtee colony ,Gokul Nagar,Tarnaka Report Status : Final Report

CLINICAL BIOCHEMISTRY

SAGE CARE 1.2

Test Name	Results	Units	Biological Reference Interval	
Iron Profile-I				
Iron(Fe) (Method: Ferrozine)	90	μg/dL	50-170	
Total Iron Binding Capacity (TIBC)	<u>454</u>	μg/dL	250-450	
Transferrin (Method: Calculated)	317.48	mg/dL	250-380	
Iron Saturation((% Transferrin Saturation)	19.82	%	15-50	
Unsaturated Iron Binding Capacity (UIBC)	364	ug/dL	110-370	

Interpretation:

- Serum transferrin (and TIBC) high, serum iron low, saturation low. Usual causes of depleted iron stores include blood loss, inadequate dietary iron. RBCs in moderately severe iron deficiency are hypochromic and microcytic. Stainable marrow iron is absent. Serum ferritin decrease is the earliest indicator of iron deficiency if inflammation is absent.
- Anemia of chronic disease: Serum transferrin (and TIBC) low to normal, serum iron low, saturation low or normal. Transferrin decreases with many inflammatory diseases. With chronic disease there is a block in movement to and utilization of iron by marrow. This leads to low serum iron and decreased erythropoiesis. Examples include acute and chronic infections, malignancy and renal failure.
- Sideroblastic Anemia: Serum transferrin (and TIBC) normal to low, serum iron normal to high, saturation high.
- Hemolytic Anemia: Serum transferrin (and TIBC) normal to low, serum iron high, saturation high.
- Hemochromatosis: Serum transferrin (and TIBC) slightly low, serum iron high, saturation very high.
- Protein depletion: Serum transferrin (and TIBC) may be low, serum iron normal or low (if patient also is iron deficient). This may occur as a result of malnutrition, liver disease, renal disease.
- Liver disease: Serum transferrin variable; with acute viral hepatitis, high along with serum iron and ferritin. With chronic liver disease (eg, cirrhosis), transferrin may be low. Patients who have cirrhosis and portacaval shunting have saturated TIBC/transferrin as well as high ferritin.

*** End Of Report ***









Page 10 of 11

Name : Mrs. S NAGAMANI

Sample ID : A1841792

Age/Gender : 60 Years/Female Reg. No : 0312503100002

Referred by : Dr. PRADEEP KUMAR MISHRA SPP Code : SPL-CV-172

Referring Customer : V CARE MEDICAL DIAGNOSTICS Collected On : 10-Mar-2025 08:46 AM
Primary Sample : Whole Blood Received On : 10-Mar-2025 01:06 PM
Sample Tested In : Serum Reported On : 10-Mar-2025 03:04 PM

Client Address : Kimtee colony ,Gokul Nagar,Tarnaka Report Status : Final Report

CLINICAL BIOCHEMISTRY

SAGE CARE 1.2

Test Name	Results	Units	Biological Reference Interval	
Thyroid Profile-I(TFT)				
` ,	88.77	ng/dL	40-181	
T3 (Triiodothyronine)		J		
T4 (Thyroxine) (Method: CLIA)	11.0	μg/dL	3.2-12.6	
TSH -Thyroid Stimulating Hormone	1.75	μIU/mL	0.35-5.5	

Pregnancy & Cord Blood

T3 (Triiodothyronine	e):	T4 (Thyroxine)	TSH (Thyroid Stimulating Hormone)
First Trimester	: 81-190 ng/dL	15 to 40 weeks:9.1-14.0 μg/dL	First Trimester : 0.24-2.99 µIU/mL
Second&Third Trimes	ster :100-260 ng/dL		Second Trimester: 0.46-2.95 µIU/mL
			Third Trimester : 0.43-2.78 μIU/mL
Cord Blood: 30-70 ng	ı/dL	Cord Blood: 7.4-13.0 μg/dL	Cord Blood: : 2.3-13.2 µIU/mL

Interpretation:

- Thyroid gland is a butterfly-shaped endocrine gland that is normally located in the lower front of the neck. The thyroid's job is to make thyroid hormones, which are secreted into the blood and then carried to every tissue in the body. Thyroid hormones help the body use energy, stay warm and keep the brain, heart, muscles, and other organs working as they should.
- Thyroid produces two major hormones: triiodothyronine (T3) and thyroxine (T4). If thyroid gland doesn't produce enough of these hormones, you may experience symptoms such as weight gain, lack of energy, and depression. This condition is called hypothyroidism.
- Thyroid gland produces too many hormones, you may experience weight loss, high levels of anxiety, tremors, and a sense of being on a high. This is called hyperthyroidism.
- TSH interacts with specific cell receptors on the thyroid cell surface and exerts two main actions. The first action is to stimulate cell reproduction and hypertrophy. Secondly, TSH stimulates the thyroid gland to synthesize and secrete T3 and T4.
- The ability to quantitate circulating levels of TSH is important in evaluating thyroid function. It is especially useful in the differential diagnosis of primary (thyroid) from secondary (pituitary) and tertiary (hypothalamus) hypothyroidism. In primary hypothyroidism, TSH levels are significantly elevated, while in secondary and tertiary hypothyroidism, TSH levels are low.







