

LABORATORY TEST REPORT

Name	: Mrs. RAJYA LAKSHMI		
Sample ID	: B2623239, B2623246		
Age/Gender	: 66 Years/Female	Reg. No	: 0312505030005
Referred by	: Dr. SRIKANTH	SPP Code	: SPL-CV-172
Referring Customer	: V CARE MEDICAL DIAGNOSTICS	Collected On	: 03-May-2025 08:05 AM
Primary Sample	: Whole Blood	Received On	: 03-May-2025 04:19 PM
Sample Tested In	: Plasma-NaF(F), Plasma-NaF(PP)	Reported On	: 03-May-2025 05:11 PM
Client Address	: Kimtee colony ,Gokul Nagar,Tarnaka	Report Status	: Final Report



CLINICAL BIOCHEMISTRY

GLUCOSE POST PRANDIAL (PP)

Test Name	Results	Units	Biological Reference Interval
-----------	---------	-------	-------------------------------

Glucose Fasting (F) **125** mg/dL 70-100
(Method: Hexokinase)

Interpretation of Plasma Glucose based on ADA guidelines 2024

Diagnosis	FastingPlasma Glucose(mg/dL)	2hrsPlasma Glucose(mg/dL)	HbA1c(%)	RBS(mg/dL)
Prediabetes	100-125	140-199	5.7-6.4	NA
Diabetes	> = 126	> = 200	> = 6.5	>=200(with symptoms)

Reference: Diabetes care 2024 Jan (1:47 (suppl.1):S20- S42.

Glucose Post Prandial (PP) **136** mg/dL 70-140
(Method: Hexokinase (HK))

Interpretation of Plasma Glucose based on ADA guidelines 2018

Diagnosis	FastingPlasma Glucose(mg/dL)	2hrsPlasma Glucose(mg/dL)	HbA1c(%)	RBS(mg/dL)
Prediabetes	100-125	140-199	5.7-6.4	NA
Diabetes	> = 126	> = 200	> = 6.5	>=200(with symptoms)

Reference: Diabetes care 2024 Jan (1:47 (suppl.1):S20- S42.

- Postprandial glucose level is a screening test for Diabetes Mellitus
- If glucose level is >140 mg/dL and <200 mg/dL, then GTT (glucose tolerance test) is advised.
- If level after 2 hours = >200 mg/dL diabetes mellitus is confirmed.
- Advise HbA1c for further evaluation.

*** End Of Report ***



Handwritten Signature
DR. LAVANYA LAGISETTY
MD BIOCHEMISTRY

LABORATORY TEST REPORT

Name	: Mrs. RAJYA LAKSHMI		
Sample ID	: B2623240		
Age/Gender	: 66 Years/Female	Reg. No	: 0312505030005
Referred by	: Dr. SRIKANTH	SPP Code	: SPL-CV-172
Referring Customer	: V CARE MEDICAL DIAGNOSTICS	Collected On	: 03-May-2025 08:05 AM
Primary Sample	: Whole Blood	Received On	: 03-May-2025 01:34 PM
Sample Tested In	: Serum	Reported On	: 03-May-2025 05:11 PM
Client Address	: Kimtee colony ,Gokul Nagar,Tarnaka	Report Status	: Final Report


CLINICAL BIOCHEMISTRY

Test Name	Results	Units	Biological Reference Interval
-----------	---------	-------	-------------------------------

Thyroid Profile-I(TFT)

T3 (Triiodothyronine) <small>(Method: CLIA)</small>	69.08	ng/dL	40-181
T4 (Thyroxine) <small>(Method: CLIA)</small>	13.79	µg/dL	3.2-12.6
TSH -Thyroid Stimulating Hormone <small>(Method: CLIA)</small>	13.37	µIU/mL	0.50-8.9

Pregnancy & Cord Blood

T3 (Triiodothyronine):	T4 (Thyroxine)	TSH (Thyroid Stimulating Hormone)
First Trimester : 81-190 ng/dL	15 to 40 weeks:9.1-14.0 µg/dL	First Trimester : 0.24-2.99 µIU/mL
Second&Third Trimester :100-260 ng/dL		Second Trimester: 0.46-2.95 µIU/mL
		Third Trimester : 0.43-2.78 µIU/mL
Cord Blood: 30-70 ng/dL	Cord Blood: 7.4-13.0 µg/dL	Cord Blood: : 2.3-13.2 µIU/mL

Interpretation:

- Thyroid gland is a butterfly-shaped endocrine gland that is normally located in the lower front of the neck. The thyroid's job is to make thyroid hormones, which are secreted into the blood and then carried to every tissue in the body. Thyroid hormones help the body use energy, stay warm and keep the brain, heart, muscles, and other organs working as they should.
- Thyroid produces two major hormones: triiodothyronine (T3) and thyroxine (T4). If thyroid gland doesn't produce enough of these hormones, you may experience symptoms such as weight gain, lack of energy, and depression. This condition is called hypothyroidism.
- Thyroid gland produces too many hormones, you may experience weight loss, high levels of anxiety, tremors, and a sense of being on a high. This is called hyperthyroidism.
- TSH interacts with specific cell receptors on the thyroid cell surface and exerts two main actions. The first action is to stimulate cell reproduction and hypertrophy. Secondly, TSH stimulates the thyroid gland to synthesize and secrete T3 and T4.
- The ability to quantitate circulating levels of TSH is important in evaluating thyroid function. It is especially useful in the differential diagnosis of primary (thyroid) from secondary (pituitary) and tertiary (hypothalamus) hypothyroidism. In primary hypothyroidism, TSH levels are significantly elevated, while in secondary and tertiary hypothyroidism, TSH levels are low.

*** End Of Report ***



*TESTS CONDUCTED @ CENTRAL LAB, HYDERABAD


 DR. LAVANYA LAGISETTY
 MD BIOCHEMISTRY

Page 2 of 2